

## 化学組成及び同位体比に基づく南鳥島周辺EEZ内堆積物の起源の変遷 Origin of deep-sea sediments within the Minamitorishima EEZ based on downhole variation of bulk chemical composition and isotopic ratios

\*田中 えりか<sup>1</sup>、安川 和孝<sup>1,2</sup>、中村 謙太郎<sup>1</sup>、宮崎 隆<sup>3</sup>、Bogdan Vaglarov<sup>3</sup>、藤永 公一郎<sup>2,1</sup>、岩森 光<sup>3,4</sup>、加藤 泰浩<sup>1,2,3</sup>

\*Erika Tanaka<sup>1</sup>, Kazutaka Yasukawa<sup>1,2</sup>, Kentaro Nakamura<sup>1</sup>, Takashi Miyazaki<sup>3</sup>, Bogdan Vaglarov<sup>3</sup>, Koichiro Fujinaga<sup>2,1</sup>, Hikaru Iwamori<sup>3,4</sup>, Yasuhiro Kato<sup>1,2,3</sup>

1. 東京大学工学系研究科、2. 千葉工業大学、3. 海洋研究開発機構、4. 東京工業大学

1. School of Engineering, The University of Tokyo, 2. Chiba Institute of Technology, 3. JAMSTEC, 4. Tokyo Institute of Technology

Recently, deep-sea sediments containing a high concentration of rare-earth elements and yttrium (REY), which is called “REY-rich mud”, were discovered in the Pacific Ocean [1]. Moreover, the presence of “extremely REY-rich mud” was confirmed within the Japanese exclusive economic zone (EEZ) surrounding Minamitorishima Island [2]. Although previous studies have shown that the main host phase of REY in these sediments is biogenic calcium phosphate based on bulk and *in situ* geochemical analyses [2,3], the contribution of other components constituting the REY-rich mud have not been completely understood yet.

One of the most effective ways to constrain the origin(s) of the deep-sea sediments, including REY-rich mud, is to use isotopic ratios of the sediments, because each possible geochemical end-member has characteristic isotopic ratios reflecting source materials and genetic processes [4].

Here, to clarify the origin of deep-sea sediments within the Minamitorishima EEZ, we modeled and quantified the contribution ratios of a few, representative geochemical end-members based on Sr isotopic ratio and major-/trace-element contents. The reconstructed downhole variation of the contribution ratios indicates a transition of end-members along with the change of geological settings due to the motion of the Pacific plate.

### References

- [1] Kato et al. (2011) *Nature Geoscience* 4, 535-539.
- [2] Iijima et al. (2016) *Geochemical Journal* 50, 557-573.
- [3] Kon et al. (2014) *Resource Geology* 64, 47-57.
- [4] Capo et al. (1998) *Geoderma* 82, 197 -225.

キーワード：Sr同位体、端成分、深海堆積物、南鳥島EEZ、レアアース泥

Keywords: Sr isotopic ratio, end-members, deep-sea sediments, Minamitorishima EEZ, REY-rich mud