

Multiple microanalysis of melt inclusions from the Pitcairn basalts and its implications for melting source region

*Takahiro Ozawa¹, Hikaru Iwamori^{1,2}, Takeshi Hanyu², Morihisa Hamada², Kenji Shimizu³, Takayuki Ushikubo³, Jun-Ichi Kimura², Qing Chang², Hitomi Nakamura^{1,2,4}, Motoo Ito³

1. Department of Earth and Planetary Sciences, Graduated School of Science and Engineering, Tokyo Institute of Technology, 2. Department of Solid Earth and Geochemistry, Japan Agency for Marine-Earth Science and Technology, 3. Kochi Core Center, Japan Agency for Marine-Earth Science and Technology, 4. ORCeNG, Chiba Institute of Technology

We have developed a systematic analytical procedure to determine 44 major-trace-volatile element concentrations and lead isotopic ratios for melt inclusions with a few tens- to hundred-micrometer in diameter. This was accomplished by using multiple microanalytical instruments including SIMS, EPMA, and LA-ICP-MS. We analyzed olivine-hosted melt inclusions in basalt lavas and a pyroclastic rock from Tedsid Volcanics, Pitcairn Island. Tedsid Volcanics represents the shielding phase of island (0.95-0.76Ma) and shows lower lead isotopic ratios than basalts from other geological formations of the post-erosional phase between 0.67 and 0.45 Ma (Pulawana Volcanics, Christians Cave Formation and Adamstown Volcanics) which show MORB-like lead isotopic ratios.

Melt inclusions in the lavas contain pargasite, ilmenite, garnet, and pyrite of up to 63 modal %. Some of the crystals could have crystallized before the melt inclusions were entrapped in the host olivine. Two groups of the melt inclusion samples were prepared for the instrumental analysis; (1) original samples, and (2) homogenized samples that were reheated at the liquidus temperature, which was carefully determined by the melting experiments, and were then quenched to produce a uniform glass. By analyzing the two suites, compositional modification during the homogenization procedure, which is commonly used in melt inclusion studies, can be discussed by comparing the analytical results of the two groups. The melt inclusions in the pyroclastic rock are naturally homogenous, with which homogenized samples were also prepared for the comparison.

Major and trace element compositions of the original and the homogenized samples overlap with the bulk compositions of their host rocks. Volatile elements are affected by the homogenization. The F contents in the homogenized samples show two trends; one exhibits a broadly constant concentration (584-842 ppm) irrespective of their various H₂O contents, whereas another shows a positive correlation with H₂O. The homogenization affects H₂O contents in the melt inclusions in several ways; increase by melting of hydrous minerals in the melt inclusions, decrease by diffusion into bubbles formed in the melt inclusions, and decrease by diffusional escape to the outside of melt inclusions. CO₂ in the melt inclusions also re-equilibrated with the bubbles formed during homogenization. The range of lead isotope ratio of melt inclusions is larger than bulk of basalt in Tedsid Volcanics. Low lead isotope ratio (²⁰⁷Pb/²⁰⁶Pb and ²⁰⁸Pb/²⁰⁶Pb) is consistent with the post-erosional phase in Pitcairn Island. The heterogeneity in Pitcairn Island is classified four major clusters based on the k-means cluster analysis. The main cluster is broadly distributed around the average lead isotopic ratios of Tedsid Volcanics. This cluster involves melt inclusions from both the basalt lavas and the pyroclastic rock. Melt inclusions from the pyroclastic rock are all glassy and appear uniform under the microscopes, yet classified into two distinct clusters; the main cluster and another cluster with a highly enriched signature. It is suggested that an enriched source that has been metasomatized by either aqueous fluid or melt and a depleted MORB-source-like material coexisted in a very small spatial scale in melting source region of the Pitcairn basalts.

