

## Reexamination of the radiometric ages of the middle Miocene near trench magmatism in SW Japan

\*新正 裕尚<sup>1</sup>、折橋 裕二<sup>2</sup>、安間 了<sup>3</sup>、角井 朝昭<sup>4</sup>、和田 穰隆<sup>5</sup>、周藤 正史<sup>6</sup>

\*Hironao Shinjoe<sup>1</sup>, Yuji Orihashi<sup>2</sup>, Ryo Anma<sup>3</sup>, Tomoaki Sumii<sup>4</sup>, Yutaka WADA<sup>5</sup>, Masafumi Sudo<sup>6</sup>

1. 東京経済大学経営学部、2. 東京大学地震研究所、3. 筑波大学生命環境系、4. 産業技術総合研究所地質調査総合センター、5. 奈良教育大学地学教室、6. ポツダム大学地球環境科学研究所

1. Faculty of Business Administration, Tokyo Keizai University, 2. Earthquake Research Institute, University of Tokyo, 3. Faculty of Life and Environmental Science, University of Tsukuba, 4. Geological Survey of Japan, National Institute of Advanced Industrial Science and Technology, 5. Department of Earth Sciences, Nara University of Education, 6. Institute of Earth and Environmental Science, University of Potsdam

In the middle Miocene time of the southwest Japan, widespread magmatism occurred in near trench region. The region of magmatism extends ca. 800 km along with SW Japan Arc, and up to 100 km in the across arc direction. It was almost contemporaneous with the clockwise rotation of SW Japan arc and the commencement of subduction of the young hence hot Shikoku Basin of the Philippine Sea plate beneath the SW Japan forearc sliver. Near trench magmatism can be divided into three types by the lithologies and the distance from the Nankai trench. Setouchi Volcanic Rocks in the Setouchi Geologic Province are composed of rhyolitic to dacitic rocks, calc-alkaline andesite including high-Mg andesite, and olivine tholeiite. Some of the felsic rocks with HREE depletion are assumed to be formed by melting of sediment on the subducting slab (Shinjoe et al., 2007). High-Mg andesite magma thought to be formed by reaction of the slab melt with mantle wedge peridotite (Tatsumi, 2006). In the Outer Zone of SW Japan are distributed Outer Zone Granitic Rocks including felsic to intermediate plutonic and volcanic rocks. Most of them are S-type felsic rocks formed by melting of deep part (ca. 20 km) of the accretionary complex (Shinjoe, 1997). Some of the igneous complexes ejected large scale ignimbrite and formed caldera. In the region closest to the Nankai trench, small igneous complexes and dikes are distributed, including the basaltic rocks both tholeiitic and alkaline affinities with or without felsic igneous rocks. Ages of these igneous rocks have been constrained mainly by conventional K-Ar and fission-track methods. The results of the dating range 10 to 17 Ma, so the igneous activities have been thought to generally contemporaneous with clockwise rotation of SW Japan Arc. However, more precise data are needed to constrain the tectonic environment of this short-lived peculiar magmatism. For example, to support the slab melting hypothesis of high-Mg andesite of the Setouchi Volcanic Rocks, magmatism should have taken place after the initiation of the subduction of the hot Shikoku Basin. We have conducted zircon U-Pb dating by LA-ICP-MS in the Earthquake Research Institute of the University of Tokyo for systematically representative igneous bodies in Kyushu, western Shikoku and Kii peninsula. Preliminary <sup>40</sup>Ar/<sup>39</sup>Ar dating was also carried out at Potsdam University for zircon-free mafic lithologies. Based on the results of dating, we demonstrate the time-space distribution of the middle Miocene near trench igneous rocks in SW Japan. The overall results of our dating are as follows. (1) U-Pb ages of the Setouchi Volcanic Rocks and Outer Zone Granitic Rocks in Kii peninsula and Kyushu island range 15.5 to 14.0 Ma, 15.1 to 12.4 Ma and 15.6 to 13.4 Ma. Thusly, along arc variation of the age of the near trench igneous rocks is not observed. (2) The oldest age samples of Outer Zone Granitic Rocks were usually obtained from the igneous bodies in the closest area to the trench, and weak across arc trend of the age variation is found in Outer Zone Granitic Rocks. (3) Comparing to the recent discussion on the completion period of the clockwise rotation of SW Japan arc (Hoshi et al., 2015; Sawada et al., 2013), all the magmatism occurred after the rotation. (4) Since the ages of the alkaline basaltic rocks with OIB-like trace element signatures also range 15.2 to 11.7 Ma, i.e. after the rotation, these magmas should have been supplied from the

mantle beneath the subducting Philippine Sea Plate.

キーワード：中期中新世、マグマティズム、ジルコンウラン・鉛年代、 $40\text{Ar}/39\text{Ar}$ 年代、四国海盆、日本海拡大

Keywords: middle Miocene, magmatism, zircon U-Pb age,  $40\text{Ar}/39\text{Ar}$  age, Shikoku Basin, Japan Sea opening