

メッセンジャー探査機のベクトル磁場データが捉えた水星の核 Mercury's core detected by MESSENGER's vector magnetic data

桂 貴暉¹、*藤 浩明²

Takaaki Katsura¹, *Hiroaki TOH²

1. 京都大学大学院理学研究科地球惑星科学専攻、2. 京都大学大学院理学研究科附属地磁気世界資料解析センター

1. Division of Earth and Planetary Sciences, Graduate School of Science, Kyoto University, 2. Data Analysis Center for Geomagnetism and Space Magnetism, Graduate School of Science, Kyoto University

MESSENGER (MErcury Surface, Space Environment, Geochemistry, and Ranging) is the first probe launched into Mercury's polar orbits and observed the planet's electromagnetic (EM) environment including its vector magnetic field from 2011 to 2015.

This study is the first to clarify the characteristic periods of magnetic field variations at Mercury and to extract the induced magnetic field explicitly from the observed data. In order to discuss EM induction within Mercury, it is necessary to acquire time-series of internal and external Gauss coefficients. However, it is very difficult to determine those coefficients as in the case of the Earth due to the limited spatial distribution of the magnetic field data of MESSENGER. For the structure of Mercury's magnetic field, low degree zonal coefficients have been determined from spherical harmonic analysis of all MESSENGER data (Anderson et al., 2012). However, the temporal variation of Mercury's magnetic field has not yet been discussed so far because the Gauss coefficients estimated in the previous study are based on the annual means. For the EM induction in Mercury, there has been only an example estimating dipolar magnetic fields induced at the top of the Mercury's core by time-varying magnetospheric field (Johnson et al., 2016). They calculated the time-varying external field from a stationary magnetic model by regarding the model as a function of the subsolar distance that shows a clear annual variation due mainly to the large eccentricity of the Mercury's orbit.

In this study, we estimated one of the most important parameters of the Mercury's internal structure, namely, its core radius by extracting time-varying internal Gauss coefficients from observed data not only for the axial dipole but also other zonal components. First, we revealed the characteristic periods of the Mercury's magnetic field variations by automatically identifying the position of the Mercury's magnetopause, converting them into the subsolar distances, and analyzing their frequency dependence. It was found that the characteristic periods that may contribute to EM induction within Mercury are mainly one Mercury year and half Mercury year. Second, we derived the time series data of internal Gauss coefficients by employing the KT14 stationary magnetic field model for Mercury (Korth et al., 2015) as a function of the subsolar distance. Finally, we estimated the core radius of Mercury using the ratio of the internal to external Gauss coefficients for the axial dipole and for one Mercury year. A stable estimate of core radius, 2019 ± 172 km, was obtained using time segments when the Mercury's magnetic field is quiet at its magnetopause. This result is within a reasonable range compared with other results determined independently from EM method such as Harder and Schubert (2001), Hauck et al. (2013) and so on.

キーワード：水星金属核、メッセンジャー探査機、ベクトル磁場データ、内外分離、年周変化、磁気圏界面
Keywords: Mercury's metallic core, MESSENGER Spacecraft, vector magnetic data, internal/external separation, annual variation, magnetopause