

Retrograde phases of former bridgmanite inclusions in superdeep diamonds?

Zedgenizov Dmitry^{1,4}, *鍵 裕之²、大谷 栄治³、辻森 樹⁴、小松 一生²、鎌田 誠司⁵
 Dmitry Zedgenizov^{1,4}, *Hiroyuki Kagi², Eiji Ohtani³, Tatsuki Tsujimori⁴, Kazuki Komatsu², Seiji Kamada⁵

1. ロシア科学アカデミーソボレフ地質学鉱物学研究所、2. 東京大学大学院理学系研究科、3. 東北大学大学院理学研究科、4. 東北大学東北アジア研究センター、5. 東北大学学際科学フロンティア研究所

1. V.S.Sobolev Institute of Geology and Mineralogy, Siberian Branch of the Russian Academy of Sciences, 2. Graduate School of Science, The University of Tokyo, 3. Graduate School of Science, Tohoku University, 4. Center for Northeast Asian Studies, Tohoku University, 5. Frontier Research Institute for Interdisciplinary Sciences, Tohoku University

Bridgmanite (Mg,Fe)SiO₃, a high pressure silicate with a perovskite structure, is dominant material in the lower mantle at the depths from 660 to 2700 km and therefore is probably the most abundant mineral in the Earth. Although synthetic analogues of this mineral have been well studied, no naturally occurring samples had ever been found in a rock on the planet's surface except in some shocked meteorites. Due to its unstable nature under ambient conditions, this phase undergoes retrograde transformation to a pyroxene-type structure. The identification of the retrograde phase as 'bridgmanite' in so-called superdeep diamonds was based on the association with ferropericlasite (Mg,Fe)O and other high-pressure (supposedly lower-mantle) minerals predicted from theoretical models and HP-HT experiments. In this study pyroxene inclusions in diamond grains from Juina (Brazil), one single-phase (Sample SL-14) and two composite inclusions of (Mg,Fe)SiO₃ coexisting with (Mg,Fe)₃Al₂Si₃O₁₂ (Sample SL-13), and with (Mg,Fe)₃Al₂Si₃O₁₂ and (Mg,Fe)₂SiO₄ (Sample SL-80) have been analyzed to identify retrograde phases of former bridgmanite. XRD and Raman spectroscopy have revealed that these are orthopyroxene (Opx). (Mg,Fe)₂SiO₄ and (Mg,Fe)₃Al₂Si₃O₁₂ in these inclusions are identified as olivine and jeffbenite (TAPP). These inclusions are associated with inclusions of (Mg,Fe)O (SL-14), CaSiO₃ (SL-80) and composite inclusion of CaSiO₃+CaTiO₃ (SL-13). XRD patterns of (Mg,Fe)SiO₃ inclusions indicate that they consist of polycrystals. This polycrystalline textures together with high lattice strain of host diamond around these inclusions observed from EBSD may be an evidence for the retrograde phase transition of former bridgmanite.

Single-phase inclusions of (Mg,Fe)SiO₃ in superdeep diamonds are suggested to represent a retrograde phase of bridgmanite and fully inherit its initial chemical composition, including a high Al and low Ni contents [Harte, Hudson, 2013; Kaminsky, 2017]. The composite inclusions of (Mg,Fe)SiO₃ with jeffbenite and other silicate and oxide phases may be interpreted as exsolution products from originally homogeneous bridgmanite [Walter et al., 2011]. The bulk compositions of these composite inclusions are rich in Al, Ti, and Fe which are similar to Al-rich bridgmanite produced in experiments on the MORB composition. However, the retrograde origin of composite inclusions due to decomposition of Al-rich bridgmanite may be doubtful because each of observed phases may represent single-phase inclusions, i.e. bridgmanite and high pressure garnet (majoritic garnet), with similar compositional features.

This work has been partially supported by RFBR (16-05-00451 and 17-55-50062).

References

Harte B., Hudson N.F.C. (2013) Mineral Associations in Diamonds from the Lowermost Upper Mantle and Uppermost Lower Mantle // D.G. Pearson et al. (eds.), Proceedings of 10th International Kimberlite Conference, Special Issue of the Journal of the Geological Society of India, Vol.1, 235-253.

Kaminsky F.V. (2017) The Earth' s Lower Mantle. Springer International Publishing AG.
Walter M.J., Kohn S.C., Araujo D., Bulanova G.P., Smith C.B., Gaillou E., Wang J., Steele A., Shirey S.B.
(2011) Deep mantle cycling of oceanic crust: evidence from diamonds and their mineral inclusions.
Science 334 (6052): 54-57.

キーワード：深部起源ダイヤモンド、ブリッジマナイト、包有物

Keywords: superdeep diamonds, bridgmanite, inclusions