

Understanding of low velocity anomaly in the asthenosphere inferred from elastic wave velocity measurement of partially molten rocks

*大橋 智典¹、坂巻 竜也¹、久野 直毅¹、肥後 祐司²、柴崎 裕樹³、鈴木 昭夫¹

*Tomonori Ohashi¹, Tatsuya Sakamaki¹, Naoki Hisano¹, Yuji Higo², Yuki Shibazaki³, Akio Suzuki¹

1. 東北大学大学院理学系研究科、2. (財)高輝度光科学センター、3. 東北大学 学際科学フロンティア研究所

1. Department of Earth Science, Tohoku University, 2. Japan Synchrotron Radiation Research Institute, 3. Frontier Research Institute for Interdisciplinary Sciences, Tohoku University

The low velocity anomaly at the top of asthenosphere is one of important seismological features, and expected to be formed by carbon dioxide (CO₂)-induced partial melting of mantle rock. For better constraint on the melt fraction and distribution in the low-velocity zone, a comparison between experimental velocity and geophysical observation is a straightforward approach. Here we have measured the elastic wave velocity of olivine + carbonate melt using in-situ ultrasonic technique under conditions relevant to the top of asthenosphere. Both of compressional (V_p) and shear (V_s) wave velocities became smaller with increasing the volume fraction of melt in the sample, and we found quantitatively the velocity attenuation as a function of melt fraction. Comparing with the low-velocity zone, we succeeded in understanding a suitable amount of melt in the asthenosphere. The presence of magmas below the lithosphere could serve as a lubricant for the dynamics of plate tectonics and be a magma source of petit-spot volcanoes located at the outer rise of the oceanic plate.

キーワード：部分溶融、弾性波速度、アセノスフェア

Keywords: partial melting, elastic wave velocity, asthenosphere