

ロシア極東サハリンから産出したメラノフロジャイトの単結晶X線回折実験

Single-crystal X-ray diffraction study of melanophlogite from Sakhalin, Far East Russia

*門馬 綱一¹、長瀬 敏郎²、ジェンキンス ロバート³、宮嶋 佑典⁴、谷 健一郎¹、井尻 暁⁶、Sergei Kasatkin⁵、Igor Chekryzhov⁵、宮脇 律郎¹

*Koichi Momma¹, Toshiro Nagase², Robert Jenkins³, Yusuke Miyajima⁴, Kenichiro Tani¹, Akira Ijiri⁶, Sergei Kasatkin⁵, Igor Chekryzhov⁵, Ritsuro Miyawaki¹

1. 独立行政法人国立科学博物館、2. 東北大学総合学術博物館、3. 金沢大学、4. 京都大学、5. ロシア科学アカデミー極東地質研究所、6. 海洋研究開発機構

1. National Museum of Nature and Science, 2. Tohoku University Museum, 3. Kanazawa University, 4. Kyoto University, 5. Far East Geological Institute, Russian Academy of Sciences, 6. JAMSTEC

Melanophlogite is one of silica clathrate minerals, which contain various gas molecules in their cage-like framework structures.

We recently discovered new localities of melanophlogite in the South Sakhalin, Far East Russia. In Nevelsk, melanophlogite occurs in calcareous concretion of the Miocene mudstone in association with autochthonous fossils of chemosynthetic fauna such as *Conchocele* sp. and *Acharax* sp. Associate minerals are opal and chalcedony, and melanophlogite occurs as aggregates of cubic crystals of sizes up to 0.1 mm.

In “Zamiraylova head” cape, melanophlogite occurs in hydrothermal veins filling fissures of a small basaltic volcanic body. The vein consists of microcrystalline quartz, opal, aragonite, calcite, and pyrite. A lot of cubic pseudomorphs of chalcedony after melanophlogite occur in voids of these veins, while unaltered melanophlogite crystals were only found at the center of thick vein. The same kind of hydrothermal veins are also distributed in a basaltic volcanic body in Kuznetsova cape.

Presence of methane in the samples from “Zamiraylova head” cape was confirmed by C-H stretching mode of the Raman spectra at $\sim 2900\text{ cm}^{-1}$. Single-crystal X-ray diffraction experiment at room temperature (23°C) revealed that its crystal structure is consistent with previously reported model, i.e., three-fold twinning of the pseudo-cubic tetragonal structure with space group $P42/nbc$ (#133). At 100°C , all reflections arising from $2\times 2\times 1$ super-cell disappeared and the structure could be refined using cubic $Pm-3n$ (#223) space group. When it was cooled down to 70°C , additional reflections reappeared at positions slightly shifted from those expected for $2\times 2\times 1$ super-cell, indicating a possible presence of long-range order or incommensurate phase between alpha-beta transition of melanophlogite.

キーワード：メラノフロジャイト、シリカクラスレート、超構造

Keywords: melanophlogite, silica clathrate, superstructure