

Mg isotopic composition of submarine vent fluids from arc and back-arc hydrothermal systems in the western Pacific

*EOM JIWON¹、荒岡 大輔²、吉村 寿紘³、蒲生 俊敬¹、川幡 穂高¹

*EOM JIWON¹, Daisuke Araoka², Toshihiro Yoshimura³, Toshitaka Gamo¹, Hodaka Kawahata¹

1. 東京大学大気海洋研究所、2. 産業技術総合研究所、3. 海洋研究開発機構

1. Atmosphere and Ocean Research Institute, The University of Tokyo, 2. AIST, 3. JAMSTEC

The major source of dissolved Mg to the ocean is continental weathering of carbonate and silicate minerals, mostly carried by rivers and groundwaters. Since only almost half of Mg from the river is buried into the sediments, hydrothermal circulation is thought to be clearly an important mechanism for Mg removal from the seawater in that Mg is completely removed during high-temperature seawater-rock interaction. However, its role in global geochemical mass balances is still highly uncertain and no research has been reported about Mg isotopes in hydrothermal solution yet. We analyzed Mg isotopic compositions of hydrothermal water samples collected from 11 sites at five arc and back-arc hydrothermal systems in the western Pacific and investigated Mg isotopic exchange between fluids and rock/sediments during hydrothermal circulation. The results showed that the Mg concentrations ranged from 0.8 mmol/kg to 46.3 mmol/kg and that the Mg isotopic ratios are from -1.18 ‰ to -0.80 ‰. Especially, sediment-starved hydrothermal solution showed that the Mg isotopic compositions tend to have lower compositions as the Mg concentrations decrease. On the other hand, sediment-hosted hydrothermal solution (e.g., CLAM site) had Mg isotopic composition of -0.81 ‰ and -0.80 ‰, slightly higher than that of seawater (-0.83 ‰). Particularly, sample which had near zero concentration of Mg (e.g. sediment-starved hydrothermal system: Suiyo Seamount) showed the lightest Mg isotopic compositions (-1.18 ‰) in our data.

キーワード：マグネシウム同位体、海底熱水

Keywords: Mg isotope, hydrothermal vent fluid