

## Stress state in the source region of M2-3 earthquakes in South African deep gold mines

\*矢部 康男<sup>1</sup>、小笠原 宏<sup>2</sup>、伊藤 高敏<sup>3</sup>、船戸 明雄<sup>4</sup>、Durrheim Raymond<sup>5</sup>、Yilmaz Halil<sup>6</sup>、Hofmann Gerhard<sup>7</sup>、阿部 周平<sup>1</sup>

\*Yasuo Yabe<sup>1</sup>, Hiroshi Ogasawara<sup>2</sup>, Takatoshi Ito<sup>3</sup>, Akio Funato<sup>4</sup>, Raymond Durrheim<sup>5</sup>, Halil Yilmaz<sup>6</sup>, Gerhard Hofmann<sup>7</sup>, Shuhei Abe<sup>1</sup>

1. 東北大学大学院理学研究科附属地震・噴火予知研究観測センター、2. 立命館大学理工学部、3. 東北大学流体科学研究所、4. 深田地質研究所、5. School of Geoscience, University of the Witwatersrand、6. School of Mining Engineering, University of the Witwatersrand、7. AngloGold Ashanti

1. Research Center for Prediction of Earthquakes and Volcanic Eruptions, Graduate School of Science, Tohoku University, 2. Faculty of Science and Engineering, Ritsumeikan University, 3. Institute of Fluid Science, Tohoku University, 4. Fukada Geological institute, 5. School of Geoscience, University of the Witwatersrand, 6. School of Mining Engineering, University of the Witwatersrand, 7. AngloGold Ashanti

As a part of the ICDP project "Drilling into seismogenic zones of M2.0 - M5.5 earthquake in South African gold mines (DSeis)", we evaluate the stress states in the source regions of Mw2.2 earthquake in Mponeng mine (M-event) and of M3.5 earthquake in Savuka mine (S-event), both of which occurred in dykes at depths of 3.3-3.5 km. A scientific borehole was drilled across the fault of the M-event and two exploration boreholes were drilled across the fault zone of the S-event. States of borehole walls (borehole breakout, BB) and of core samples (core discing, CD) recovered from the boreholes suddenly changed across the fault or the fault zones, indicating a significant contrast between the stress states in the hanging-wall and the footwall of the source faults. Two stress measurement techniques (DRA and DCDA) were applied to the core samples recovered from the three boreholes. Differential stresses were found to be higher in the dyke compared with those in the host rock. Combining the stresses measured by these techniques with the occurrence criteria of BB and CD, the principal stress state in the source region of the M-event was estimated. The vertical stress was as high as ~180 MPa in the dyke in the hanging-wall of the fault, while it was ~100 MPa in the dyke in the footwall. This contrast between the vertical stresses in two sides of the fault can be attributed to the stress redistribution by an inhomogeneous slip on a non-planar fault, which is consistent with the pre-seismic stress distribution on the fault calculated by a numerical modeling of the stress perturbations due to nearby mining. The stress state in the source region of S-event significantly differed from the virgin stress state in this mine district. This should be due to the stress disturbance associated with nearby mining, as well as to the stress redistribution induced by the S-event. However, its spatial variation was too complicated to be explained by a simple model as in the case of the M-event.

キーワード : Mining induced earthquake、南アフリカ大深度金鉱山、震源域の応力場、ICDP DSeis

Keywords: Mining induced earthquake, South African deep mine, Stress state in the source region, ICDP DSeis