

Postseismic fluid discharge chemically recorded in altered pseudotachylyte discovered from an ancient megasplay fault

*長谷川 亮太¹、山口 飛鳥¹、福地 里菜²、濱田 洋平³、北村 有迅⁴、木村 学⁵、芦 寿一郎¹、石川 剛志³

*Hasegawa Ryota¹, Asuka Yamaguchi¹, Rina Fukuchi², Yohei Hamada³, Yujin Kitamura⁴, Gaku Kimura⁵, Juichiro Ashi¹, Tsuyoshi Ishikawa³

1. 東京大学大気海洋研究所、2. 海洋研究開発機構地震津波海域観測研究開発センター、3. 海洋研究開発機構高知コア研究所、4. 鹿児島大学大学院理工学研究科、5. 東京海洋大学海洋資源環境学部

1. Atmosphere and Ocean Research Institute, The University of Tokyo, 2. Research and Development (R&D) Center for Earthquake and Tsunami, Japan Agency for Marine-Earth Science and Technology, 3. Kochi Institute for Core Sample Research, Japan Agency for Marine-Earth Science and Technology, 4. Graduate School of Science and Engineering KAGOSHIMA UNIVERSITY, 5. Tokyo University of Marine Science and Technology School of Marine Resources and Environment

Megasplay fault branching from plate boundaries at subduction zones are thought to be important source of earthquakes generating tsunami. In this study, we performed structural and geochemical analyses on a fossilized Megasplay fault (the Nobeoka Thrust of the Shimanto accretionary complex) to understand fluid-rock interaction and how the splay fault plays a role of fluid flow in the seismogenic zone.

Nobeoka Thrust is a low-angle thrust which subdivides the Shimanto belt in Kyushu into the northern (Cretaceous and Tertiary) and the southern (Tertiary) subbelts, and is an exhumed analogue of an ancient megasplay fault. The hanging wall and the footwall of the Nobeoka Thrust show difference in lithology and metamorphic grade, and their maximum burial temperature is estimated from vitrinite reflectance analysis to be 320~330°C and 250~270°C, respectively.

As a result of structural observation, the principal slip zone (PSZ) of the Nobeoka thrust is composed of foliated cataclasite originated from sandstone-shale mélange including thin (~1.5 mm thick) pseudotachylyte layer. Major and trace element composition analysis and EPMA element mapping revealed that the pseudotachylyte within the PSZ is enriched in Li and Cs, as well as slip zone of a minor fault in the footwall. Li- and Cs-enrichment in pseudotachylyte is interpreted as a result of fluid-rock interaction just after faulting (postseismic stage) because such an anomaly is only formed by large fluid/rock ratio ($R > 512$ to 24 at 250 to 350 degrees C) under the existence of Li- and Cs-enriched fluid. X-ray diffraction analysis showed that the pseudotachylyte was devitrified to form palygorskite and muscovite, similar to the pseudotachylyte founded in minor shear zone of the hanging wall (Okamoto et al., 2006). The amount of fluid reacted with pseudotachylyte is estimated to be 4.8×10^0 to 1.64×10^4 m³, based on empirical relationships among fault thickness, displacement and fault length.

キーワード：流体－岩石相互作用、シュードタキライト、巨大分岐断層

Keywords: fluid-rock interaction, pseudotachylyte, megasplay fault