

火道を上昇するマグマ内における水の減圧発泡に関する理論的研究 -粘性項を含む核形成速度の導出と気泡数密度の再評価-

A new formulation of viscosity-included bubble nucleation rate in ascending magmas and re-estimation of bubble number density

*西脇 瑞紀¹、寅丸 敦志²

*Mizuki Nishiwaki¹, Atsushi Toramaru²

1. 九州大学大学院理学府地球惑星科学専攻、2. 九州大学大学院理学研究院地球惑星科学部門

1. Department of Earth and Planetary Sciences, Graduate School of Sciences, Kyushu University, 2. Department of Earth and Planetary Sciences, Faculty of Sciences, Kyushu University

The dynamics in the conduit controls the surface phenomena such as the shift between explosive and nonexplosive eruptions. Therefore, quantifying the unobservable behavior of magmas in the conduits during eruptions is useful as gaining a much better understanding of eruptions. In particular, the formation process of bubbles plays an important role as the driving force for eruptions. There have been various kinds of studies to quantitatively understand the formation process of bubbles in magmas, e.g. numerical simulations, laboratory experiments, and textural analysis of bubbles. Last one allows us to estimate the decompression rate of the ascending magma in the conduit. Specifically, BND (Bubble Number Density) in pyroclasts such as pumice or scoria has been focused and utilized as the good index. In the previous works, the viscosity effects of silicate melts only have been taken into account in the growth process incorporating the Rayleigh-Plesset equation as the momentum equation. On the other hand, the nucleation rate is assumed to be independent of viscosity, since the formula is derived by only considering the diffusional migration of the gas molecules in liquid phase to gas phase, neglecting the change of the gas molar volume. However, in the case of the vesiculation in the high-viscous melt, it can be intuitively expected that the nucleation becomes difficult, because viscous flow in the melt surrounding an expanding bubble can work as the powerful resistance against bubble nucleation. Thus, we have to formulate the nucleation rate taking into account the viscosity effects of liquid. According to Kagan's method for the bubble nucleation in a single component system, we have derived the nucleation rate for a two component system (silicate-water system). The nucleation rate explicitly includes the viscosity term in the pre-exponential factor in terms of Péclet number. Péclet number is the ratio of two timescales: a timescale of the bubble growth by diffusion and a timescale of viscous relaxation. So Péclet number is inversely proportional to viscosity. In the previous works the viscosity term is not included in the pre-exponential factor, namely the situation where Péclet number is positive infinity, is considered. On the other hand, in this study, the factor drastically decreases by orders of magnitude as the value of Péclet number decreases below unity under a given supersaturation. Values of Péclet number calculated from real physical properties of magmas are in this range, so existing formulae of nucleation rates lead to overestimated rates.

We have numerically solved the time evolution of bubble nucleation and growth in magmas decompressed in a constant rate by using the newly derived nucleation rate formula. Like similar to the results obtained in the previous version, two regimes have been found on the relationship among BND (the value when a magma reached at the surface), diffusivity and viscosity. They are: the diffusion-controlled regime where BND does not depend on viscosity but does only on diffusivity, and the viscosity-controlled regime where BND exponentially increases with the increase of viscosity but independent of diffusivity. Compared with the previous study, the transition point of two regimes moves to the higher-viscosity region by a factor of order unity and BND values are dramatically reduced by several

orders of magnitude in the viscosity-controlled regime. In addition, as viscosity increases beyond the transition point, BND increases by several orders of magnitude with increasing viscosity by one order, taking the maximum value at the certain value of viscosity, and then diminishes to zero at infinite viscosity. This behavior is consistent with our intuition that no nucleation occurs in extremely viscous magmas. These results can be expected to account for laboratory experimental results which are partly inconsistent with the previous BND model.

キーワード：気泡核形成、気泡成長、核形成速度、気泡数密度、ペクレ数、粘性

Keywords: bubble nucleation, bubble growth, nucleation rate, BND (Bubble Number Density), Péclet number, viscosity