

T-PARCIIのドロップゾンデ観測と気象庁全球予測システムを用いた台風 Trami(2018)を対象とする観測システム実験

Observing System Experiment for Typhoon Trami (2018) using T-PARCII dropsondes and JMA Global Forecasting System

*山口 宗彦¹、石橋 俊之¹、中澤 哲夫¹、伊藤 耕介²、山田 広幸²、大東 忠保³、長浜 則夫⁴、清水 健作⁴、久保田 尚之⁵、高橋 幸弘⁵、加藤 雅也⁶、金田 幸恵⁶、吉岡 真由美⁶、篠田 太郎⁶、高橋 暢宏⁶、坪木 和久⁶

*Munehiko Yamaguchi¹, Toshiyuki Ishibashi¹, Tetsuo Nakazawa¹, Kosuke Ito², Hiroyuki Yamada², Tadayasu Ohigashi³, Norio Nagahama⁴, Kensaku Shimizu⁴, Hisayuki Kubota⁵, Yukihiro Takahashi⁵, Masaya Kato⁶, Sachie Kanada⁶, Mayumi Yoshioka⁶, Taro Shinoda⁶, Nobuhiro Takahashi⁶, Kazuhisa Tsuboki⁶

1. 気象庁気象研究所、2. 琉球大学理学部、3. 防災科学技術研究所、4. 明星電気株式会社、5. 北海道大学大学院理学院、6. 名古屋大学宇宙地球環境研究所

1. Meteorological Research Institute, Japan Meteorological Agency, 2. Faculty of Science, University of the Ryukyus, 3. National Research Institute for Earth Science and Disaster Resilience, 4. Meisei Electric, 5. Graduate School of Science, Hokkaido University, 6. Institute for Space-Earth Environmental Research, Nagoya University

Airborne typhoon observations were conducted for Typhoon Trami (2018) under Tropical cyclones-Pacific Asian Research Campaign for Improvement of Intensity estimations/forecasts (T-PARCII). Using the dropsonde data (zonal and meridional winds, temperature, and relative humidity) and the Japan Meteorological Agency Global Spectral Model (JMA/GSM, TL959L100) and 4 Dimensional Variational data assimilation (4DVAR, TL319L100), we performed Observing System Experiment (OSE) to see the impact of the newly available observational data on the track/intensity predictions by the JMA/GSM.

Three sets of numerical experiments, CNTL, EXP1 and EXP2, were performed. All the T-PARCII dropsonde and typhoon bogus data were used in the data assimilation in CNTL while T-PARCII dropsonde (typhoon bogus) data only were assimilated in the 4DVAR in EXP1 (EXP2). The period of the experiments is from 2018.09.25 00UTC to 2018.09.28 18UTC with a 6-hour data assimilation and forecasting cycle.

The error of the track predictions of EXP1 and EXP2 is smaller than that of CNTL. For example, the 3-day track prediction error of CNTL, EXP1 and EXP2 is 232, 201 and 218 km, respectively. A possible reason of the degradation of the track predictions by CNTL is that some dropsonde data were assimilated at the almost same locations as typhoon bogus data, which might deteriorate the initial structure of the typhoon.

In the operational configuration of the 4DVAR, all dropsonde data were assimilated irrespective of the existence of the typhoon bogus data. The results of a series of these numerical experiments may lead to improvement in the operational setting of dropsonde assimilation.

キーワード：航空機観測、台風、気象予測

Keywords: Airborne observation, Typhoon, Weather Prediction