

微小重力下における毛管中の水分移動速度の低下

The slowing down of water movement in capillary tubes under microgravity.

*丸尾 裕一¹、佐藤 直人¹、長沼 菜摘¹、野川 健人¹、登尾 浩助²

*Yuichi Maruo¹, Naoto Sato¹, Natsumi Naganuma¹, Kento Nogawa¹, Kosuke Noborio²

1. 明治大学大学院、2. 明治大学

1. Graduate school of Agriculture, Meiji University, 2. Meiji University

The international deep space gateway is going to be built on the moon orbit from early 2020s. As manned space missions may get farther and longer, the importance of food production in space increases. For food production by growing crops in space, water by the crops plays an important role; however, few workers have conducted research on water movement in porous media under microgravity. It was reported that a water infiltration rate in porous media was slower under microgravity than under 1G. The objective of this study is to reveal what makes water move slower in porous media under microgravity. Capillary rises in capillary tubes, which were simpler pores, were observed under microgravity induced by a free-fall tower. The measured rates of capillary rise were slower than theoretical values calculated by Lucas-Washburn equation for 1G condition. The slower rates could be explained if either surface tension, contact angle, or viscosity are the function of gravity. Previous research reported that larger viscosity was observed under microgravity than under 1G although surface tension and contact angle were not changed by gravity. Slower water flows in capillary tubes under microgravity may be explained by changes in viscosity as a function of gravity.

キーワード：土壌物理学、微小重力、水分移動

Keywords: Soil physics, Microgravity, Water movement