

## Gas origin of hydrate-bound gas in Lake Baikal

\*Jumpei Matsuda<sup>1</sup>, Yuki Kikuchi<sup>1</sup>, Akihiro Hachikubo<sup>1</sup>, Oleg Khlystov<sup>2</sup>, Gennadiy Kalmychkov<sup>3</sup>, Marc De Batist<sup>4</sup>, Hirotohi Sakagami<sup>1</sup>, Hirotsugu Minami<sup>1</sup>, Satoshi Yamashita<sup>1</sup>

1. Kitami Institute of Technology, 2. Limnological Institute, SB RAS, 3. Vinogradov Institute of Geochemistry, SB RAS, 4. Ghent University

In the framework of international collaboration between Kitami Institute of Technology (Japan), Limnological Institute (Russia), and Ghent University (Belgium), natural gas hydrates have been recovered from lake-bottom sediments at Lake Baikal. Near-surface gas hydrate was first discovered at the Malenky mud volcano at the southern Baikal basin in 2000. Multi-phase Gas Hydrate Project (MHP, 2009-2018) has revealed distribution of near-surface gas hydrates at the southern and central Baikal basins. We found six new places (Enkhelook, Soukhaya, KIT, LIN, PosolCanyon1, and PosolCanyon2) during the VER18-03 cruise in 2018. The total number of places where near-surface gas hydrates were retrieved is 53. In this report, we summarize characteristics of hydrate-bound gases retrieved from all the sites since 2005.

Gas hydrate crystals were recovered from lake-bottom sediment cores, and hydrate-bound gas samples were stored in 5-mL vials. Total number of hydrate-bound gas samples was 626. Molecular and isotopic compositions of hydrate-bound gas were measured using a gas chromatograph and a stable isotope ratio mass spectrometer.

In the Bernard plot,  $C_1 \delta^{13}C$  and  $C_1 / (C_2 + C_3)$  distributes from -70‰ to -44‰ and from 6 to 100,000, respectively, suggesting that the origin of hydrate-bound gas is microbial, thermogenic, and their mixed-gas. Almost all data are plotted on the mixing line between microbial and thermogenic, however, Gorevoy Utes (oil seep site) showed thermogenic  $C_1$  with small composition of  $C_2$  and  $C_3$ . The data concentrates in the microbial field ( $C_1 \delta^{13}C$ : -66‰ and  $C_1 / (C_2 + C_3)$ : 1,000-10,000).  $C_1 \delta D$  of thermogenic  $C_1$  increases with  $C_1 \delta^{13}C$ .  $C_2 \delta^{13}C$  distributes from -69‰ to -23‰. 17 sites (ex. Solzan, Talanka, Turka, KIT, etc.) showed microbial  $C_1$  and  $C_2$ . 27 sites (Malenky, Oblom, Seep13, Kukuy K6, etc.) showed microbial  $C_1$  and thermogenic  $C_2$ , indicating that small amount of thermogenic gas is mixed with microbial gas. Other sites (Mamay, PosolBank, Gorevoy Utes, Kedr, etc.) showed thermogenic  $C_1$  and  $C_2$ . The Milkov plot (relation between  $C_1 \delta^{13}C$  and  $C_2 \delta^{13}C$ ) can express mixing condition between microbial and thermogenic rather than the Bernard plot.  $C_2 \delta D$  decreased with decrease in  $C_2 \delta^{13}C$ , therefore microbial  $C_2$  is both depleted in  $^{13}C$  and D. We conclude that hydrate-bound gas at Lake Baikal is mainly microbial gas, and thermogenic gas ascends from deep layer at some areas (Kedr, PosolBank, Kukuy, and Gorevoy Utes) and forms the structure II gas hydrate.

Keywords: gas hydrate, Lake Baikal, gas origin