

## Visible Photometric, Spectroscopic and Polarimetric observations of an inactive comet, P/2016 BA14 with the 1.6m Pirka Telescope

\*Ryo Yamagata<sup>1</sup>, Tomohiko Sekiguchi<sup>1</sup>, Ryo Okazaki<sup>1</sup>, Kazuya Doi<sup>1</sup>, Daisuke Kuroda<sup>2</sup>, Masateru Ishiguro<sup>3</sup>, Hiroyuki Naito<sup>4</sup>

1. Hokkaido University of Education, 2. Kyoto University, 3. Seoul National University, 4. Nayoro Observatory

We performed photometric, spectroscopic and polarimetric observations of an inactive comet, P/2016 BA14 with the 1.6m Pirka Telescope, Hokkaido.

P/2016 BA14 ( $a=3.02$  au,  $e=0.67$ ,  $i=18.9$  deg.) was discovered as an unusual asteroid with a cometary orbit by the PANSTAARS on 2016 January 21. Since Knight et al. 2016 detected its cometary tail when the object close to sun ( $r=1.018$  au). It was classified into a Jupiter Family Comets. It can be regarded as a bare nucleus invisible coma because very weak comet activity that is different from general comet. There is a remarkable similarity between the orbits of 252P/LINEAR ( $a=3.05$  au,  $e=0.67$ ,  $i=10.4$  deg) and P/2016 BA14. Jupiter Tisserand parameter is 252P/LINEAR and that of P/2016 BA14 are 2.82. P/2016 BA14 and 252P/LINEAR is division comet that is originally same comet (Jian-Yang.2017).

In this study, surface reflection properties of P/2016 BA14 classify D-type asteroids (Bus-DeMeo taxonomy) from result of visible spectroscopy (435-820nm). Color of surface is little red as D-type asteroid, but very blue as comet nucleus from result of BVRI photometry. The linear polarization  $Pr=10.46.\pm 0.11\%$  ( $\alpha=46^\circ$ ) is in close accord with that of D-type asteroids and comet nucleus.

Keywords: P/2016 BA14, Photometric, Spectroscopic, Polarimetric, D-type