

フィリピンパナイ島ヒルギダマシ属林における細根生産力の評価 Estimation of fine root productivity of mangroves based on ingrowth core method on *Avicennia* forest in Panay Island, Philippines

*小野 賢二¹、諏訪 鍊平²、レネ ローロン³、アリエル ブランコ³、ギア アルバノ³、グリーン クルツ³、ヤスミン プリマヴェーラーチロール⁴、ローナルド マリオ⁴、灘岡 和夫⁵
*ONO Kenji¹, Rempei SUWA², Rene ROLLON³, Ariel C. BLANCO³, Giannina MARIE G. ALBANO³, Green ANN A. CRUZ³, Yasumin H. PRIMAVERA-TIROL⁴, Ronald J. MALIO⁴, Kazuo NADAOKA⁵

1. 国立研究開発法人森林研究・整備機構森林総合研究所、2. 国立研究開発法人 国際農林水産業研究センター、3. フィリピン大学、4. ア克蘭州立大学、5. 東京工業大学

1. Forestry and Forest Products Research Institute, 2. Japan International Research Center for Agricultural Science, 3. University of the Philippines, 4. Aklan State University, 5. Tokyo Institute of Technology

To accurately evaluate the storing and sequestering processes of carbon of mangrove ecosystems in tropical and subtropical regions, it is necessary to determine the fine root productivity in mangrove forests. It is because high productivity of mangrove fine root highly contributes the huge carbon storage of mangrove ecosystem, especially their substratum. In the present study, we aimed to estimate the fine root productivity of *Avicennia* forest in KII eco-park in northern region of Panay Island, Philippines.

Ingrowth core experiment was conducted according to Fujimaki et al. (2004 PLSO) to estimate the annual production rate of fine root. Two transects (125 m in length) were set at each interior and seaside positions in KII eco-park, where have a different tidal frequency. Moreover, three plots were set up at respective transects and each twelve mesh cores, i.e., ingrowth cores were put on the mangrove forest floor in one plot at February, 2018. 4 cores had been already collected from the plots at September, 2018, picked up all fine roots from cores, determined their masses, and estimated their annual production rates.

Annual fine root production of *Avicennia* forest in KII eco-park were estimated to be $6.1 \pm 2.8 \text{ Mg ha}^{-1} \text{ year}^{-1}$ at interior and $7.1 \pm 6.3 \text{ Mg ha}^{-1} \text{ year}^{-1}$ at seaside respectively, which were based on the data of a half-year incubation cores.

These results indicate that FRP in mangrove forests was generally equal and/or relatively high compared with values previously recorded for *Avicennia* forests (Poungparn et al. 2012, 2015).

キーワード：細根生産量、ヒルギダマシ属、フィリピンパナイ島

Keywords: fine root productivity, *Avicennia*, Panay Island