

Abundant ferromanganese microparticles in oxic pelagic sediments

*Go-Ichiro Uramoto^{1,2}, Yuki Morono², Naotaka Tomioka², Shigeyuki Wakaki², Ryoichi Nakada², Rota Wagai³, Kentaro Uesugi⁴, Akihisa Takeuchi⁴, Masato Hoshino⁴, Yoshio Suzuki^{4,5}, Satoshi Mitsunobu⁶, Fumito Shiraishi⁷, Hiroki Suga⁵, Yasuo Takeichi⁸, Yoshio Takahashi⁵, Fumio Inagaki²

1. Kochi University, 2. Japan Agency for Marine-Earth Science and Technology, 3. National Agriculture and Food Research Organization, 4. Japan Synchrotron Radiation Research Institute, 5. University of Tokyo, 6. Ehime University, 7. Hiroshima University, 8. High Energy Accelerator Research Organization

Manganese is the third most abundant metallic element, after iron and titanium, in the Earth's crust, and ferromanganese minerals are sensitive to changes in redox condition. Assessing the formation and preservation of ferromanganese minerals is important for understanding the global marine cycles of manganese and numerous associated trace elements. The most extensive manganese mineral deposits occur on abyssal plains, including those below open-ocean gyres. However, the extent of ferromanganese minerals buried in subseafloor sediments remains unclear. During the Integrated Ocean Drilling Program (IODP) Expedition 329, we drilled the entire sedimentary sequence at 6 sites in the ultra-oligotrophic region of the South Pacific Gyre (SPG), where dissolved O₂ and aerobic microbial communities are present from the seafloor to the sediment-basement interface [1]. By using high-resolution imaging techniques, we found abundant (10⁸-10⁹ particles cm⁻³) micrometer-scale ferromanganese mineral particles (Mn-microparticles) in the oxic pelagic clays of the South Pacific Gyre [2]. Furthermore, we established microparticle separation technique by the combined use of density concentration and flowcytometer/particles sorting techniques. Major and trace element compositional analyses revealed that iron and manganese were the major components of Mn-microparticles, while they also contained rare-earth elements. From the number of Mn-microparticles per unit volume of sediment and the global distribution of oxic pelagic clays, we estimated the number of Mn-microparticles to be 1.5-8.8 × 10²⁸ particles, which accounted for 1.28-7.62 Tt of manganese. This estimate is at least 2 orders of magnitude larger than the manganese budget for manganese nodules and manganese crusts on the seafloor. Along with the examination of manganese input, formation, and preservation, this study provides new insights into the global budget of metallic elements in the abundant Mn-microparticles present in deep subseafloor environments.

[1] D' Hondt, S., Inagaki, F., Zarikian, C. A., Abrams, L. J., Dubois, N., Engelhardt, T., Evans, H., Ferdelman, T., Gribsholt, B., Harris, R. N., Hoppie, B. W., Hyun, J.-H., Kallmeyer, J., Kim, J., Lynch, J. E., McKinley, C. C., Mitsunobu, S., Morono, Y., Murray, R. W., Pockalny, R., Sauvage, J., Shimono, T., Shiraishi, F., Smith, D. C., Smith-Duque, C., Spivack, A. J., Steinsbu, B. O., Suzuki, Y., Szpak, M., Toffin, L., Uramoto, G., Yamaguchi, T. Y., Zhang, G., Zhang, X.-H., and Ziebis, W. Presence of oxygen and aerobic communities from seafloor to basement in deep-sea sediment. *Nature Geosciences*, 8(4), 299-304, 2015.

[2] Uramoto, G.-I., Morono, Y., Tomioka, N., Wakaki, S., Wagai, R., Uesugi, K., Takeuchi, A., Hoshino, M., Suzuki, Y., Shiraishi, F., Mitsunobu, S., Suga, H., Nakada, R., Takeichi, Y., Takahashi, Y., and Inagaki, F., Significant contribution of subseafloor microparticles to the global manganese budget. *Nature Communications*. **10**, 400, <https://doi.org/10.1038/s41467-019-08347-2>, 2019

Keywords: Mn-microparticle, oxic pelagic clay, IODP Exp. 329