

## 過去の海洋魚類群集を復元するための堆積物中魚類起源DNA抽出法の改善に関する研究

### Improvements of extraction methods of sedimentary fish-origin DNA for reconstruction of marine fish communities

\*倉田 純<sup>1</sup>、玉井 大道<sup>2</sup>、土居 秀幸<sup>3</sup>、源 利文<sup>4</sup>、加 三千宣<sup>5</sup>

\*Jun Kurata<sup>1</sup>, Hiromichi Tamai<sup>2</sup>, Hideyuki Doi<sup>3</sup>, Toshihumi Minamoto<sup>4</sup>, Michinobu Kuwae<sup>5</sup>

1. 愛媛大学理学部地球科学科、2. 愛媛大学大学院理工学研究科、3. 兵庫県立大学大学院シミュレーション学研究科、4. 神戸大学大学院人間発達環境学研究科、5. 愛媛大学沿岸環境科学センター

1. department of earth science, Ehime university, 2. Graduate School of Science and Engineering, Ehime university, 3. Graduate School of Simulation Studies, University of Hyogo, 4. Graduate School of Human Development and Environment, Kobe University, 5. Center for Marine Environmental Studies, Ehime university

Environmental DNA is DNA fragments derived from body fluids or skin fragments of macro organisms contained in water or soil. It is expected to can be a technology to reveal long-term fish community changes which still remains unknown in the world ocean. However, DNA in the sediments has a low concentration, therefore, it is difficult to detect for a various kind of fish species. The purpose of this research is to elucidate the method that can take sedimentary DNA more efficiently. At this time, we tested five DNA extraction methods including the conventional one together with beads beating time. Quantitative PCA results showed that Power Soil had the best yield in the replications. In addition, DNA extraction of *Trachurus japonicus* figured out a better yield by increasing the amount of sediment sample. There was a significant difference between 10 min. and 30 sec. of beads beating time, it is possible that there is a higher yield with short time beating. Our future works is to test whether DNA yield improves by shortening or eliminating bead beating, effectively removal of PCR inhibitors, and an utility of PCR inhibitor.

キーワード：環境DNA、海洋堆積物

Keywords: Environmental DNA, ocean sediment