

$^{13}\text{CH}_4$ を包接するメタンハイドレートの平衡圧

Equilibrium pressure of clathrate hydrates encaged $^{13}\text{CH}_4$

*菊池 優樹¹、八久保 晶弘¹、小関 貴弘¹、竹谷 敏²

*Yuki Kikuchi¹, Akihiro Hachikubo¹, Takahiro Ozeki¹, Satoshi Takeya²

1. 北見工業大学、2. 産業技術総合研究所

1. Kitami Institute of Technology, 2. National Institute of Advanced Industrial Science and Technology (AIST)

Methane molecule is composed of carbon and hydrogen, and three kinds of isotopologues, $^{12}\text{CH}_4$ (98.9%), $^{13}\text{CH}_4$ (1.1%), and CH_3D (0.013%) exist in nature. We often measure stable carbon isotope ($^{13}\text{C}/^{12}\text{C}$) of methane to understand its gas origin. Since their weight of isotopologues differ with each other, physicochemical properties of them are also different. Pure methane hydrate must be "mixed-gas hydrate" of their isotopologues. Ozeki *et al.* (2018) measured an equilibrium pressure of CH_3D hydrate, but that of $^{13}\text{CH}_4$ has not been studied yet. Fractionation of hydrogen isotope of methane during the formation of methane hydrate has been reported by Hachikubo *et al.* (2007) that δD of hydrate-bound methane becomes several ‰ smaller than that of residual methane. This result suggests that the equilibrium pressure of CH_3D hydrate is larger than that of CH_4 hydrate, and Ozeki *et al.* (2018) demonstrated the difference between these equilibrium pressures. Since Hachikubo *et al.* (2007) showed no isotopic fractionation in methane $\delta^{13}\text{C}$, the equilibrium pressures of $^{12}\text{CH}_4$ and $^{13}\text{CH}_4$ hydrates are thought to be almost the same. In this study, we measured the equilibrium pressures of $^{13}\text{CH}_4$ hydrate to check the difference from that of $^{12}\text{CH}_4$ hydrate.

Methane hydrate samples were synthesized in small pressure cells (volume: 5 mL). Fine ice powder (1g) was put in a pressure cell, and introduced $^{13}\text{CH}_4$ (purity: 99.5%, Taiyo-Nissan). Clathrate hydrate was formed by melting the ice powder at the temperature of 273.2K under high pressure of methane. We also prepared normal methane (purity: 99.99% for methane, but 98.9% for $^{12}\text{CH}_4$, Takachiho Chemical Industry Co. Ltd.) hydrate as a reference, using the same preparation method. These pressure cells were placed in a temperature-controlled liquid bath, and measured their equilibrium pressures from 269.5K to 277.9K.

The difference in equilibrium pressure between $^{13}\text{CH}_4$ and normal methane (mainly $^{12}\text{CH}_4$) hydrates was smaller than the measurement error. This results agree with the previous report by Hachikubo *et al.* (2007).

References

Ozeki T, Kikuchi Y, Takeya S, Hachikubo A (2018) Phase equilibrium of isotopologue methane hydrates enclathrated CH_3D and CD_4 . J Chem Eng Data 63(6): 2266-2270, doi: 0.1021/acs.jced.8b00203

Hachikubo A, Kosaka T, Kida M, Krylov A, Sakagami H, Minami H, Takahashi N, Shoji H (2007) Isotopic fractionation of methane and ethane hydrates between gas and hydrate phases. Geophys Res Lett 34: L21502. doi:10.1029/2007GL030557

キーワード：ガスハイドレート、メタンハイドレート、平衡圧、安定同位体

Keywords: gas hydrate, methane hydrate, equilibrium pressure, stable isotope