

TW Hyaの原始惑星系円盤に付随する局所的なサブミリ波超過の検出

Discovery of a localized excess in the millimeter emission of the protoplanetary disk around TW Hya

*塚越 崇¹、武藤 恭之²、野村 英子³、川邊 良平^{1,4,5}、金川 和弘⁹、奥住 聡³、井田 茂¹⁰、Walsh Catherine⁶、Millar Tom⁷、高橋 実道^{2,1}、橋本 淳⁸、鶴山 大智⁵、田村 元秀^{5,8}

*Takashi Tsukagoshi¹, Takayuki Muto², Hideko Nomura³, Ryohei Kawabe^{1,4,5}, Kazuhiro Kanagawa⁹, Satoshi Okuzumi³, Shigeru Ida¹⁰, Catherine Walsh⁶, Tom J Millar⁷, Sanemichi Takahashi^{2,1}, Jun Hashimoto⁸, Taichi Uyama⁵, Motohide Tamura^{5,8}

1. 国立天文台電波研究部、2. 工学院大学教育推進機構、3. 東京工業大学地球惑星科学系、4. 総合研究大学院大学、5. 東京大学理学系研究科、6. School of Physics and Astronomy, University of Leeds、7. Astrophysics Research Centre, Queen's University Belfast、8. 自然科学研究機構アストロバイオロジーセンター、9. 東京大学ビッグバン宇宙国際研究センター、10. 東京工業大学地球生命研究所

1. Division of Radio Astronomy, National Astronomical Observatory of Japan, 2. Division of Liberal Arts, Kogakuin University, 3. Department of Earth and Planetary Sciences, Tokyo Institute of Technology, 4. The Graduate University for Advanced Studies, 5. School of Science, University of Tokyo, 6. School of Physics and Astronomy, University of Leeds, 7. Astrophysics Research Centre, Queen's University Belfast, 8. Astrobiology Center, National Institutes of Natural Sciences, 9. Research Center for the Early Universe, University of Tokyo, 10. Earth-Life Science Institute, Tokyo Institute of Technology

Planets are formed from dust and gas in circumstellar disks around young stars.

Observations of dust and gas in circumstellar disks can provide fundamental information on planet formation processes, and high-resolution (sub-)millimeter imaging is now revealing the dust substructure likely generated by forming planets in disks.

Here we report the detection of an excess in dust continuum emission at 233 GHz (1.3 mm in wavelength) in the circumstellar disk around TW Hya, the closest such system to Earth, revealed through high-sensitivity observations with the Atacama Large Millimeter/submillimeter Array (ALMA).

We have carried out 3 au resolution continuum observations at 233 GHz (Band6) toward the protoplanetary disk (PPD) around TW Hya using ALMA in 2016 and 2017.

The sensitivity has been improved by a factor of 3 than that of our previous cycle 3 observations.

The overall structure is axisymmetric, and there are apparent gaps at 25 and 41 au as previously reported.

The most remarkable new finding is a few au scale excess emission at the south-west part of the PPD.

The excess emission is located at 52 au from the disk center and is 1.5 times brighter than the surrounding PPD.

The extracted emission after subtracting the axisymmetric PPD emission has a size of 4.4x1.0 au and a total flux density of 250 micro-Jy, corresponding to a dust mass of 0.03 earth masses.

Since the excess emission can also be marginally identified in the Band 7 image at almost the same position, the blob is unlikely a background source.

The excess emission can be explained by a massive CPD around a Neptune mass forming-planet or a dust clump accumulated in a small elongated vortex.

キーワード：惑星系形成、原始惑星系円盤、アルマ

Keywords: Planet formation, Protoplanetary disk, ALMA