

相対論的電子降り込みに対する南北両半球大気のレスポンス

Atmospheric responses in both hemispheres to relativistic electron precipitation

*田中 良昌^{1,13,14}、西山 尚典^{1,14}、門倉 昭^{1,13,14}、尾崎 光紀²、三好 由純³、大山 伸一郎^{3,15}、片岡 龍峰^{1,14}、堤 雅基^{1,14}、西村 耕司^{1,13,14}、佐藤 薫⁴、笠原 禎也²、熊本 篤志⁵、土屋 史紀⁵、吹澤 瑞貴⁵、疋島 充⁶、松田 昇也⁶、松岡 彩子⁶、篠原 育⁶、能勢 正仁³、長妻 努⁷、篠原 学⁸、藤本 晶子⁹、寺本 万里子³、野村 麗子¹⁰、行松 彰^{1,14}、細川 敬祐¹¹、小路 真史³、Ralph Latteck¹²

*Yoshimasa Tanaka^{1,13,14}, Takanori Nishiyama^{1,14}, Akira Kadokura^{1,13,14}, Mitsunori Ozaki², Yoshizumi Miyoshi³, Shin-ichiro Oyama^{3,15}, Ryuho Kataoka^{1,14}, Masaki Tsutsumi^{1,14}, Koji Nishimura^{1,13,14}, Kaoru Sato⁴, Yoshiya Kasahara², Atsushi Kumamoto⁵, Fuminori Tsuchiya⁵, Mizuki Fukizawa⁵, Mitsuru Hikishima⁶, Shoya Matsuda⁶, Ayako Matsuoka⁶, Iku Shinohara⁶, Masahito Nose³, Tsutomu Nagatsuma⁷, Manabu Shinohara⁸, Akiko Fujimoto⁹, Mariko Teramoto³, Reiko Nomura¹⁰, Akira Sessai Yukimatu^{1,14}, Keisuke Hosokawa¹¹, Masafumi Shoji³, Ralph Latteck¹²

1. 国立極地研究所、2. 金沢大学理工研究域、3. 名古屋大学宇宙地球環境研究所、4. 東京大学大学院理学系研究科、5. 東北大学大学院理学研究科、6. 宇宙航空研究開発機構宇宙科学研究所、7. 情報通信研究機構、8. 鹿児島工業高等専門学校、9. 九州工業大学、10. 国立天文台、11. 電気通信大学大学院情報理工学研究科、12. Leibniz-Institute of Atmospheric Physics e.V. at the University Rostock, Germany、13. 情報・システム研究機構 データサイエンス共同利用基盤施設 極域環境データサイエンスセンター、14. 総合研究大学院大学、15. University of Oulu, Finland

1. National Institute of Polar Research, 2. Institute of Science and Engineering, Kanazawa University, 3. Institute for Space-Earth Environmental Research, Nagoya University, 4. Graduate School of Science, The University of Tokyo, 5. Graduate School of Science, Tohoku University, 6. Japan aerospace exploration agency, Institute of space and astronautical science, 7. National Institute of Information and Communications Technology, 8. Kagoshima National College of Technology, 9. Kyushu Institute of Technology, 10. National Observatory of Japan, 11. Department of Communication Engineering and Informatics, University of Electro-Communications, 12. Leibniz-Institute of Atmospheric Physics e.V. at the University Rostock, Germany, 13. Polar Environment Data Science Center, Joint-Support Center for Data Science Research, Research Organization of Information and Systems, 14. The Graduate University for Advanced Studies (SOKENDAI), 15. University of Oulu, Finland

Relativistic electron precipitation (REP) in the polar atmosphere has been paid attention to because it changes atmospheric composition, causing ozone depletion after migrating to the lower altitude during polar winter. A plausible mechanism for REP is the interaction with magnetospheric plasma waves, such as electromagnetic ion cyclotron (EMIC) waves and whistler-mode chorus waves. Previous studies have already shown that EMIC waves can effectively drive REP, however, observational evidence that chorus waves can drive REP was lacking. In this study we present direct comparison between magnetospheric plasma waves and polar mesosphere winter echoes (PMWE) simultaneously observed with the Arase satellite and high-power Mesosphere, Stratosphere and Troposphere (MST) radars in both hemispheres, PANSY and MAARSY. PMWE can be related to the ionization in the mesosphere due to REP. The PMWE were observed at 03-07 UT on March 21, 2017 during the passage of corotating interaction region (CIR), during which the Arase footprints were close to Syowa Station (SYO; -69.00S, 39.58E), Antarctica, and Husafell (HUS; 65.67N, -21.03E), Iceland.

The observational results are summarized as follows. (1) EMIC waves and whistler-mode chorus waves were observed by Arase near the equatorial magnetosphere during 02:30-04:45 UT and 04:45-07:00 UT, respectively. (2) PMWE were detected with both the MAARSY radar at Andøya (AND; 69.30N, 16.04E), Norway, and the PANSY radar at SYO, Antarctica, during 04:45-07:00 UT, which was the recovery phase of an isolated substorm. We believe this is the first time PMWE have been observed in both hemispheres

at exactly the same time. (3) During 04:45-07:00 UT, the temporal variation of the chorus wave power was similar to those of the PMWE power in the both hemispheres. (4) The PMWE observed at SYO during 03:00-04:00 UT before the substorm onset was consistent with the occurrence of the EMIC waves. The item (2) shows direct evidence that chorus waves during the substorm recovery phase cause REP. We estimated the resonance energy of electrons interacting with the observed LBC waves, however, the estimated energy was too low to cause PMWE at an altitude lower than 70 km. However, we found that the resonance energy becomes greater than 1 MeV, if LBC waves propagate to the magnetic latitude greater than 30 degrees and resonate with energetic electrons there. As for item (4), the PMWE observed at SYO during 03:00-04:00 can be related to wave-particle interaction with EMIC waves, which may be generated inside the plasmopause by ring-current hot ions with temperature anisotropy. This anisotropy was caused by magnetospheric compression due to increasing solar wind dynamic pressure during 01:00-06:00 UT just after the arrival of the CIR. Since CIRs are main sources of geomagnetic disturbances during the declining phase of the solar cycle and solar minimum, this event is not rare but rather a common atmospheric response caused by interaction between recurrent large-scale solar wind structures and geospace.

キーワード：極域中間圏冬季エコー、あらせ衛星、PANSYレーダー、MAARSYレーダー、相対論的電子降り込み、磁気圏プラズマ波動

Keywords: Polar Mesosphere Winter Echoes, Arase satellite, PANSY radar, MAARSY radar, relativistic electron precipitation, magnetospheric plasma waves