

## An extreme erosion of the plasmasphere during the 7–10 September 2017 storm

## An extreme erosion of the plasmasphere during the 7–10 September 2017 storm

\*尾花 由紀<sup>1</sup>、丸山 奈緒美<sup>2,3</sup>、新堀 淳樹<sup>4</sup>、橋本 久美子<sup>5</sup>、Fedrizzi Mariangel<sup>2,3</sup>、能勢 正仁<sup>4</sup>、大塚 雄一<sup>4</sup>、西谷 望<sup>4</sup>、堀 智昭<sup>4</sup>、熊本 篤志<sup>6</sup>、土屋 史紀<sup>6</sup>、松田 昇也<sup>7</sup>、松岡 彩子<sup>7</sup>、笠原 禎也<sup>8</sup>、吉川 顕正<sup>9</sup>、三好 由純<sup>4</sup>、篠原 育<sup>7</sup>

\*Yuki Obana<sup>1</sup>, Naomi Maruyama<sup>2,3</sup>, Atsuki Shinbori<sup>4</sup>, Kumiko K. Hashimoto<sup>5</sup>, Mariangel Fedrizzi<sup>2,3</sup>, Masahito Nose<sup>4</sup>, Yuichi Otsuka<sup>4</sup>, Nozomu Nishitani<sup>4</sup>, Tomoaki Hori<sup>4</sup>, Atsushi Kumamoto<sup>6</sup>, Fuminori Tsuchiya<sup>6</sup>, Shoya Matsuda<sup>7</sup>, Ayako Matsuoka<sup>7</sup>, Yoshiya Kasahara<sup>8</sup>, Akimasa Yoshikawa<sup>9</sup>, Yoshizumi Miyoshi<sup>4</sup>, Iku Shinohara<sup>7</sup>

1. 大阪電気通信大学、2. University of Colorado Boulder、3. NOAA Space Weather Prediction Center、4. 名古屋大学、5. 吉備国際大学、6. 東北大学、7. 宇宙航空研究開発機構、8. 金沢大学、9. 九州大学

1. Osaka Electro-Communication University, 2. University of Colorado Boulder, 3. NOAA Space Weather Prediction Center, 4. Nagoya University, 5. Kibi International University, 6. Tohoku University, 7. Japan Aerospace Exploration Agency, 8. Kanazawa University, 9. Kyushu University

We report an extreme erosion of the plasmasphere arising from the September 2017 storm. The cold electron density is identified from the upper limit frequency of UHR waves observed by the HFA of the PWE instrument onboard the Arase spacecraft. The HFA provides the spectrograms of electric fields in a frequency range from 10 kHz to 10 MHz with an 8-sec time resolution (Kasahara et al., 2018; Kumamoto et al., 2018). Such wide frequency coverage allows observation of the UHR frequency even deep inside the plasmasphere.

The electron density profiles reveal that the plasmasphere was severely eroded during the recovery phase of the storm and the plasmopause was located at  $L \sim 1.7$  at 23 UT on 8 Sep 2017. This is the first report of such deep erosion of the plasmasphere ( $L_{pp} < 2$ ) with the in-situ observation of the electron density. The degree of the severity is much more than what is expected from the SYM-H minimum ( $-146$  nT) of the event.

We further study absolute GNSS-TEC, ground-based magnetic field, the doppler velocity obtained from the SuperDARN Hokkaido Radars to investigate the location of the midlatitude ionospheric trough, the penetration of the convection electric field to the equator and middle latitudes, respectively. Our results suggest that the middle latitude electric field was penetrated from the high-latitude storm time convection for several hours. Furthermore, the Ionosphere-Plasmasphere-Electrodynamics (IPE) model (Sun et al., 2015; Maruyama et al., 2016) is used to understand the dynamics of the plasmasphere. The IPE results suggest that the middle latitude electric field penetrates from the high-latitude storm time convection that lasts for several hours can explain the degree of severity during the September 7-11 2017 storm.

キーワード：プラズマ圏、内部磁気圏、磁気圏電離圏結合

Keywords: plasmasphere, inner magnetosphere, Magnetosphere-ionosphere coupling