

東赤道太平洋 IODP Site U1335 堆積物の古地磁気学的研究 — 過去800万年間の相対古地磁気強度と伏角異常

Paleomagnetic study of the IODP Site U1335 sediments in the eastern equatorial Pacific - relative paleointensity and inclination anomaly over the last 8 Myr

*山本 裕二¹、山崎 俊嗣²

*Yuhji Yamamoto¹, Toshitsugu Yamazaki²

1. 高知大学 海洋コア総合研究センター、2. 東京大学大気海洋研究所

1. Center for Advanced Marine Core Research, Kochi University, 2. Atmosphere and Ocean Research Institute, University of Tokyo

Obtaining continuous records of relative paleointensity (RPI) and inclination anomaly (ΔI) is inevitable to understand the fundamentals of the geodynamo, but available records older than ~ 3 Ma are still very limited in time and space. We performed a paleomagnetic study of the Integrated Ocean Drilling Program (IODP) Site U1335 sediments in the eastern equatorial Pacific to obtain continuous RPI and ΔI records since ~ 8 Ma. Slow deposition, ~ 8.4 m/Myr or less, limits the resolution of the records but did allow for determination of long-term variations. Rock-magnetic measurements showed that biogenic magnetite dominates the magnetic mineral assemblages, and the proportion of biogenic to terrigenous magnetic minerals increases prior to ~ 4 Ma. The average paleointensity between ~ 4 and 8 Ma is approximately 30% lower than that from 0 to ~ 4 Ma. The apparent reduction of RPI at ~ 4 Ma reaches approximately $\sim 50\%$, but $\sim 20\%$ of this is estimated to be artificial, induced by the increase in the proportion of biogenic magnetite. No relation between paleointensity and polarity length is recognized for the last ~ 8 Myr. The magnitude of ΔI is slightly larger during reversed polarity chrons ($4.43^\circ \pm 1.47^\circ$) than normal polarity chrons ($0.69^\circ \pm 2.98^\circ$) over the last ~ 5 Myr, which agrees with the available TAF models of this time span. Prior to ~ 6 Ma, the sign of ΔI during the normal chrons might have switched to positive, and ΔI during reversed chrons might have been slightly larger than that after ~ 5 Ma.