## 2016年熊本地震震源域の応力場の時空間変化 Spatiotemporal variations of stress field in the 2016 Kumamoto earthquake (M 7.3) area

\*趙 大鵬<sup>1</sup>、Yu Zhiteng<sup>2</sup>、Li Jiabiao<sup>2</sup>、Huang Zhouchuan<sup>3</sup> \*Dapeng Zhao<sup>1</sup>, Zhiteng Yu<sup>2</sup>, Jiabiao Li<sup>2</sup>, Zhouchuan Huang<sup>3</sup>

1. 東北大学大学院理学研究科附属地震・噴火予知研究観測センター、2. 中国海洋局第二海洋研究所、3. 南京大学地球科学 系

1. Department of Geophysics, Tohoku University, 2. State Oceanic Administration, 3. School of Earth Sciences and Engineering, Nanjing University

Focal mechanism solutions of 349 local crustal earthquakes and stress tensors in central-north Kyushu are determined using a large number of P-wave polarity data and a 3-D velocity model. The obtained tensional ( $\sigma$  3) axis trends NNW-SSE or N-S horizontally, and the compressive ( $\sigma$  1) axis trends WSW-ENE or E-W. The direction of the  $\sigma$ 3 axis is consistent with the spreading direction at the Okinawa Trough. The stress field in the 2016 Kumamoto earthquake area is attributed to rollback of the Philippine Sea plate and northward extension of the Okinawa Trough. The orientation of the  $\sigma$ 3 axis is quite stable, whereas the  $\sigma$ 1 and  $\sigma$ 2 axes are distributed in a wide range. With the 3-D velocity model, the principal stress axes can be better determined by inverting a large number of FMSs. Spatial and temporal variations of the stress field are well revealed in the Kumamoto source area. The  $\sigma 3$  axis significantly rotated counterclockwise after the M 6.5 foreshock, and rotated clockwise after the M 7.3 mainshock on the Futagawa fault segment. The stress rotation suggests a small magnitude of deviatoric stress in the source area, which indicates a small friction coefficient on the seismogenic faults. In addition, a generally small value of friction (~0.4) on the faults is obtained by the stress inversions, indicating that the faults in the Kumamoto earthquake area are weak. The fault weakening may be caused by the arc magma and fluids ascending from the mantle wedge associated with the dehydration reactions of the subducting Philippine Sea slab.

キーワード: 熊本地震、地殻応力、断層メカニズム Keywords: Kumamoto earthquake, crustal stress, focal mechanism