## Greenhouse gases and air quality monitoring system for urban anthropogenic emission estimation around Jakarta megacity Greenhouse gases and air quality monitoring system for urban anthropogenic emission estimation around Jakarta megacity

\*西橋 政秀<sup>1</sup>、向井 人史<sup>1</sup>、寺尾 有希夫<sup>1</sup>、橋本 茂<sup>1</sup>、Rizaldi Boer<sup>2</sup>、Muhammad Ardiansyah<sup>2</sup>

、Bregas Budianto<sup>2</sup>、Adi Rakhman<sup>2</sup>、Gito Sugih Immanuel<sup>2</sup>、Rudi Nugroho<sup>3</sup>、Nawa Suwedi<sup>3</sup> 、Anies Marufatin<sup>3</sup>、Muhammad Agus Salim<sup>3</sup>、Dodo Gunawan<sup>4</sup>、Eka Suharguniyawan<sup>4</sup>、Asep Firman Ilahi<sup>4</sup>、Muharam Syam Nugraha<sup>4</sup>、Ronald Christian Wattimena<sup>4</sup>、Bayu Feriaji<sup>4</sup> 、Qoriana Maulani<sup>4</sup>、小田 知宏<sup>5</sup>、Thomas Lauvaux<sup>6</sup>

\*Masahide Nishihashi<sup>1</sup>, Hitoshi Mukai<sup>1</sup>, Yukio Terao<sup>1</sup>, Shigeru Hashimoto<sup>1</sup>, Rizaldi Boer<sup>2</sup>, Muhammad Ardiansyah<sup>2</sup>, Bregas Budianto<sup>2</sup>, Adi Rakhman<sup>2</sup>, Gito Sugih Immanuel<sup>2</sup>, Rudi Nugroho <sup>3</sup>, Nawa Suwedi<sup>3</sup>, Anies Marufatin<sup>3</sup>, Muhammad Agus Salim<sup>3</sup>, Dodo Gunawan<sup>4</sup>, Eka Suharguniyawan<sup>4</sup>, Asep Firman Ilahi<sup>4</sup>, Muharam Syam Nugraha<sup>4</sup>, Ronald Christian Wattimena<sup>4</sup>, Bayu Feriaji<sup>4</sup>, Qoriana Maulani<sup>4</sup>, Tomohiro Oda<sup>5</sup>, Thomas Lauvaux<sup>6</sup>

1. 国立環境研究所、2. ボゴール農科大学、3. インドネシア技術評価応用庁、4. インドネシア気象気候地球物理庁、5. USRA/NASA、6. LSCE

1. National Institute for Environmental Studies, 2. IPB University, Indonesia, 3. Agency for the Assessment and Application of Technology (BPPT), Indonesia, 4. Meteorological, Climatological, and Geophysical Agency (BMKG), Indonesia, 5. Universities Space Research Association/NASA Goddard Space Flight Center, USA, 6. Laboratory for Sciences of Climate and Environment (LSCE), France

National Institute for Environmental Studies (NIES) has implemented an international joint research project for greenhouse gases (GHGs) and air quality monitoring with three institutes in Indonesia, IPB, BPPT, and BMKG since 2015/2016. The purpose of this project is to quantify anthropogenic emissions from Jakarta megacity and characterize them in terms of socioeconomic activities in the city. In order to respond to the Paris Climate Agreement, it is important for a monitoring project like ours not only to be capable of monitoring the increasing anthropogenic emissions by rapid economic growth in a developing country, but also to assess future those reduction impacts resulted from mitigation strategies implemented. It is also important to observe GHGs and related air pollutants with high accuracy in Indonesia because such kinds of observations are very limited in Southeast Asia.

We have maintained continuous monitoring systems of CO<sub>2</sub>, CH<sub>4</sub>, CO, NO<sub>x</sub>, SO<sub>2</sub>, O<sub>3</sub>, aerosol concentrations (PM<sub>2.5</sub>, PM<sub>10</sub>, BC) and the chemical components (NO<sub>3</sub><sup>-</sup>, SO<sub>4</sub><sup>-2</sup>) of PM<sub>2.5</sub> and PM<sub>10</sub>, and meteorological parameters at three sites: Serpong (Jakarta suburb), Bogor (center of Bogor city), and Cibeureum (mountainous area, background-like site) since 2016/2017. We have also performed automatic flask sampling of ambient air once a week. The air samples are used to analyze N<sub>2</sub>O, SF<sub>6</sub>, and carbon isotopes (<sup>13</sup>C, <sup>14</sup>C) in CO<sub>2</sub> at NIES and to validate CO<sub>2</sub>, CH<sub>4</sub>, and CO data obtained from the continuous measurement.

We have also conducted high-resolution atmospheric  $CO_2$  simulations using the Weather Research and Forecasting model coupled to Chemistry (WRF-Chem). We used two emission inventories to prescribe the surface emissions: ODIAC (Open-source Data Inventory for Anthropogenic  $CO_2$ ) as fossil fuel  $CO_2$  (ff $CO_2$ ) and MsTMIP (Multi-scale Synthesis and Terrestrial Model Intercomparison Project) as biogenic  $CO_2$ 

## (bioCO<sub>2</sub>).

We analyzed the intersite differences of daytime  $CO_2$  mole fractions ( $dCO_2$ ) between the urban sites (Serpong, Bogor) and the background-like site (Cibeureum) in the dry season (July-August 2017) and the rainy season (January-February 2018). The observed  $dCO_2$  at Serpong was 8.7 and 0.9 ppm in the dry and rainy seasons, respectively. The simulated  $dCO_2$  at Serpong shows similar decreasing trend, which is 8.5 and 3.1 ppm in the dry and rainy season possibly due to the poor reproducibility of meteorological fields (wind environment near the surface), the simulation indicates two main factors of the seasonal differences in  $dCO_2$ : one is the reduction of  $ffCO_2$  at Serpong in the rainy season (3.3 ppm) and another one is the reduction of  $bioCO_2$  uptake by photosynthesis at Cibeureum in the rainy season (2.9 ppm). The seasonal differences in  $dCO_2$  observed and simulated at Bogor are similar to those of Serpong, but smaller. In our presentation, we will also present the relationship between  $CO_2$  and the other species.

キーワード:温室効果ガス、大気質、都市観測、CO2シミュレーション、インドネシア Keywords: greenhouse gases, air quality, urban monitoring, CO2 simulation, Indonesia