## Extension of log *K* of aqueous species to low-water density conditions and comparison with hydrothermal experiments on feldspar dissolution

## \*石井 肇<sup>1</sup>、岡本 敦<sup>1</sup>、大柳 良介<sup>2</sup>、土屋 範芳<sup>1</sup> \*Hajime Ishii<sup>1</sup>, Atsushi Okamoto<sup>1</sup>, Ryosuke Oyanagi<sup>2</sup>, Noriyoshi Tsuchiya<sup>1</sup>

1. 東北大学大学院環境科学研究科、2. 国立研究開発法人海洋研究開発機構

1. Graduate School of Environmental Studies, Tohoku University, 2. Japan Agency for Marine-earth Science and Technology

In order to obtain the thermodynamic properties of aqueous species (ions and complexes) at supercritical conditions, the Helgeson-Kirham-Flowers (HKF) electrostatic model has been widely used. However, that model cannot be applied to the region close to critical point of water and low-water density region (< 0.35 g/cm<sup>3</sup>). In this study, we extended the logarithm of equilibrium constant (log K) data in SUPCRT92 with slop07.dat thermodynamic dataset to the vapor-like supercritical region using the density model instead of the HKF model. We extrapolated log K for 21 aqueous species and 45 representative minerals in the system of Si-Al-Ca-Na-K-H<sub>2</sub>O-HCl-SO<sub>4</sub>. For validating the extended log K data, we conducted the hydrothermal experiments for dissolution of feldspar (albite powder, K-feldspar-albite-quartz powder) at temperatures of 400, 420 °C, and pressures of 20, 25, 30, 35 MPa, respectively. We compared the extrapolated log K with the ion products (log Q) obtained from the solution chemistry of the experiments and extrapolated log K data of aqueous species. The difference between log K and log Q was almost less than 2 in logarithm unit, which is lower than that extrapolated by the HKF model in SUPCRT92. These results suggest that the density-based extended log K may be useful as the first approximation at temperatures of > 400 °C. In contrast, in vapor region at lower temperature of < 400 °C, the extrapolation is not acceptable due to large density gap at the vapor-saturated curve. Now, we have conducted the flow-through hydrothermal experiments on mineral precipitation in low-water density region using an input solution basalt- and granite-dissolved solutions. In contrast to quartz and metastable silica phases precipitated from the granite-dissolve solutions, from the basalt-dissolved solutions platy minerals and prism minerals precipitated, which are probably albite and clay minerals. We compare the experimental results with the reaction path modeling expanded log K data, and reveals the differences in granite-hosted and basalt-hosted hydrothermal systems.