

Forearc Basin Separation, Inner Wedge Structure, and Megathrust Segmentation of the Nankai Forearc, Japan

*木村 学¹、中村 恭之¹、白石 和也¹、藤江 剛¹、小平 秀一¹、辻 健²、福地 里菜³、山口 飛鳥⁴

*Gaku Kimura¹, Yasuyuki Nakamura¹, Kazuya Shiraishi¹, Gou Fujie¹, Shuichi Kodaira¹, Takeshi Tsuji², Rina Fukuchi³, Asuka Yamaguchi⁴

1. 国立研究開発法人海洋研究開発機構、2. 九州大学、3. 鳴門教育大学、4. 東京大学大気海洋研究所

1. Japan Agency for Marine-Earth Science and Technology, 2. Kyushu University, 3. Naruto University of Education, 4. Atmosphere and Ocean Research Institute, The University of Tokyo

The Nankai forearc, Japan, consists of inner and outer wedges with cover sediments. The forearc basin is separated into five subbasins based on topography: Enshu, Kumano, Muroto, Tosa, and Hyuga. The separation is linked to the rupture area of large earthquakes. Structural examination of the forearc basin and inner wedge off the Kii Peninsula suggests that thick cover sediments and underlying accretionary prism thrust on the landward hard basement of the igneous complex off Cape Shionomisaki. Uplift of the inner wedge due to oblique back-thrusting caused separation of the Kumano and Muroto forearc basins. The igneous complex basement is situated on the plate boundary megathrust of the hypocenters pertaining to the 1944 Tonankai and 1946 Nankai earthquakes. The denser rocks than the surrounding accretionary complex of the inner wedge might have worked as an upper plate keystone on the seismogenic megathrust and separated the rupture area of both the earthquakes. The geological heterogeneity of the upper plate partly contributes to the heterogeneous distribution of shallow, very low-frequency earthquakes. The geological composition of the Nankai forearc is a significant clue for revealing the forearc dynamics of subduction zones in general.

キーワード：付加体、南海トラフ、前弧海盆、地震発生帯、沈み込み帯、プレート境界

Keywords: accretionary wedge, Nankai trough, forearc basin, seismogenic zone, subduction zone, plate boundary