## A modeling study of effective radiative forcing and climate response due to increased methane concentration

\*Bing Xie<sup>1,2</sup>, Hua Zhang<sup>1,2</sup>, Dong-dong Yang<sup>3</sup>, Zhi-li Wang<sup>4</sup>

1. Laboratory for Climate Studies of China Meteorological Administration, National Climate Center, China Meteorological Administration, 2. Collaborative Innovation Center on Forecast and Evaluation of Meteorological Disasters, Nanjing University of Information Science & Technology, 3. College of Atmospheric Science, Nanjing University of Information Science & Technology, 4. State Key Laboratory of Severe Weather & Key Laboratory of Atmospheric Chemistry of CMA, Chinese Academy of Meteorological Sciences

An atmospheric general circulation model BCC\_AGCM2.0 and observation data from ARIS were used to calculate the effective radiative forcing (ERF) due to increased methane concentration since pre-industrial times and its impacts on climate. The ERF of methane from 1750 to 2011 was 0.46 W m<sup>-2</sup> by taking it as a well-mixed greenhouse gas, and the inhomogeneity of methane increased its ERF by about 0.02 W m<sup>-2</sup>. The change of methane concentration since pre-industrial led to an increase of 0.31 °C in global mean surface air temperature and 0.02 mm d<sup>-1</sup> in global mean precipitation. The warming was prominent over the middle and high latitudes of the Northern Hemisphere (with a maximum increase exceeding 1.4 °C). The precipitation notably increased (maximum increase of 1.8 mm d<sup>-1</sup>) over the ocean between 10°N and 20°N and significantly decreased (maximum decrease >–0.6 mm d<sup>-1</sup>) between 10°S and 10°N. These changes caused a northward movement of precipitation cell in the Intertropical Convergence Zone (ITCZ). Cloud cover significantly increased (by approximately 4%) in the high latitudes in both hemispheres, and sharply decreased (by approximately 3%) in tropical areas.

Keywords: Methane, Effective radiative forcing, Climate change