

## Climatological variations of tropospheric CO<sub>2</sub> over the Asia-Pacific region observed by the CONTRAIL commercial airliner measurements

\*Taku Umezawa<sup>1</sup>, Hidekazu Matsueda<sup>2</sup>, Yosuke Niwa<sup>2</sup>, Yousuke Sawa<sup>2</sup>, Toshinobu Machida<sup>1</sup>, Lingxi Zhou<sup>3</sup>

1. National Institute of Environmental Studies, 2. Meteorological Research Institute, 3. Chinese Academy of Meteorological Sciences

We present spatial and temporal variations of tropospheric CO<sub>2</sub> over the Asia-Pacific region analyzed from 10 years of the CONTRAIL commercial airliner measurements. The CONTRAIL flights with the high-frequency CO<sub>2</sub> measurements have covered large part of the Asia-Pacific region. Here we address climatological variations of CO<sub>2</sub> from the boundary layer to the upper troposphere across the Asia-Pacific region toward comprehensive understanding of influence of Asian surface fluxes under the varying seasonal meteorology. Highlights of this study are summarized as follows. (1) Seasonally elevated and highly variable CO<sub>2</sub> is observed in East Asia to the North Pacific in spring. This is likely explained by active passage of eastward-tracking synoptic systems that sweeps the continental East Asia and uplifts the region's CO<sub>2</sub> emissions up to the free troposphere. (2) The region-wide CO<sub>2</sub> decrease is obvious across the Asia-Pacific region and it is principally composed of CO<sub>2</sub> drawdowns originating in two distinct regions: boreal Eurasia and South Asia. We observed seasonally largest variability of CO<sub>2</sub> in the UT north of 40° N, likely due to heterogeneous spatial distributions of biospheric fluxes combined with sporadic convective vertical transport over the continent. Our observations also characterized distinct CO<sub>2</sub> depletion in the upper troposphere over South Asia as a result of strong South Asian biospheric uptakes and confinement in the Asian summer monsoon anticyclone. The development and decay of the anticyclone remarkably contributes to distributing CO<sub>2</sub> over the Asia-Pacific region. (3) As the cyclonic activity decays and the monsoon anticyclone develops from spring to summer, CONTRAIL measurements with highest data density over East Asia to the North Pacific serve as better constraints to CO<sub>2</sub> fluxes in East Asia in spring and those in South Asia in late summer.

Keywords: Carbon dioxide, Aircraft observation, Asian monsoon