

別府湾および大槌湾の浅海域の一次生産速度に地下水湧出が及ぼす影響 Influence of submarine groundwater discharge on phytoplankton primary productivity at nearshore coasts in Beppu Bay and Otsuchi Bay

*本田 尚美¹、杉本 亮²、三島 壮智³、大沢 信二³、小路 淳⁴、富永 修²、谷口 真人¹

*Hisami Honda¹, Ryo Sugimoto², Taketoshi Mishima³, Shinji Ohsawa³, Jun Shoji⁴, Osamu Tominaga², Makoto Taniguchi¹

1. 総合地球環境学研究所、2. 福井県立大学海洋生物資源学部、3. 京都大学大学院理学研究科付属地球熱学研究施設、4. 広島大学大学院生物圏科学研究科

1. Research Institute for Humanity and Nature, 2. Faculty of Marine Bioscience, Fukui Prefectural University, 3. Institute for Geothermal Sciences Graduate School of Sciences, Kyoto University, 4. Hiroshima University

In recent years, a number of studies have shown that submarine groundwater discharge (SGD) is an alternative nutrient pathway and can drive primary production in coastal seas. However, little is known about a relationship between input of nutrients through the groundwater and response of primary production. To clarify the relationship, we conducted *in situ* measurements of primary productivity (PP) using stable ¹³C tracer method under different strength sites of SGD at nearshore coasts in Beppu Bay and Otsuchi Bay during the summer in 2016. Considering the differences of light intensity and water temperature at each site, we have also incubated the bottles taken from each site under same conditions of light and water temperature on land. In both bays, significant positive relationships between *in situ* PP and *ex situ* PP ($r > 0.91$, $p < 0.01$) indicated that *in situ* PP would be controlled by nutrient availability. In Beppu Bay, *in situ* PP and ²²²Rn activity ranged from 4.4 to 23.3 $\mu\text{g C L}^{-1} \text{h}^{-1}$ and 69.8 to 586.8 Bq m^{-3} , respectively. Although there was no clear relationship between *in situ* PP and ²²²Rn activity, biomass specific PP (P_B , $\mu\text{g C } \mu\text{g chl a}^{-1} \text{ h}^{-1}$) showed the positive correlation with ²²²Rn activity. In Otsuchi Bay, *in situ* PP and ²²²Rn activity ranged from 4.5 to 10.7 $\mu\text{g C L}^{-1} \text{ h}^{-1}$ and 298 to 765.8 Bq m^{-3} , respectively. ²²²Rn activity did not related to *in situ* PP and P_B . This could be due to low phosphate concentrations in terrestrial confined groundwater. Our experimental studies suggested that the mechanism by which SGD affects phytoplankton production differs from one ecosystem to another.

キーワード：一次生産速度、ラドン222、海底湧水、浅海域

Keywords: Primary productivity, 222Rn, Submarine groundwater discharge, Nearshore coast