

Changes in main components of the water cycle of Lake Khanka during the 1949 - 2015 period.

Vladimir Georgievskiy¹, Nina Speranskaya¹, Tatiana Fuksova¹, *Pavel Groisman²

1. State Hydrological Institute, St. Petersburg, Russia, 2. Hydrology Science and Services Corporation, Asheville, North Carolina, USA

Lake Khanka (or Xingkai in China) located on the border between Primorsky Krai (Russia) and Heilongjiang Province (Northeast China) belongs to the Ussuri River basin. The Lake's drainage basin is fed by 28 rivers (8 in China and 20 in Russia), but the only outflow of the Lake is the Sungach River. During the past 15 years (2000-2015) level of the Lake have increased by 1.87 m and exceeded the historical maximum during the period of observations since 1949. It resulted in flooding of river-bank areas and agricultural lands. To understand reasons of these level changes it is necessary to analyze effects of natural and anthropogenic factors that influence the Lake water budget.

During this study of the Lake water balance, we estimated contribution of different hydrometeorological variables on the changes of its level. For these analyses we used regular hydrometeorological observations within the Lake basin from 1949 to 2015. Anthropogenic effects were estimated from the water management information and from statistical and satellite data.

Analysis of the Lake level observations showed an upward tendency since the 1980s, which became most disastrous since 2000. These changes in the Lake level were accompanied by increase of the surface water inflow into the Lake. Furthermore, since 1975, increase of annual precipitation on the Lake surface has been observed. Together, with a wind speed decrease over the Lake surface during the entire observation period, evaporation from the Lake surface has also decreased.

Anthropogenic impact on the Lake Khanka level is less evident, if compare to natural processes. In particular, during 2000-2015 the effect of natural processes on the Lake Khanka level has been more significant than of anthropogenic factors (Table). Specifically, the total surface water inflow into the Lake, precipitation at the Lake surface, and evaporation from its surface determine the lake level changes by 153 cm (or 82 % of the total change). Anthropogenic factors explain 34 cm or 18% of the Lake level changes. Among them, the effect of the Muling River flood water diversion on the Chinese territory was more important than other anthropogenic factors.

Keywords: Lake Khanka, water budget, climatic and anthropogenic changes

Table. Contribution of natural and anthropogenic factors to the Lake Khanka level changes during the 2000-2015 period.

Factors	Contribution to the Lake level changes	
	in cm	in % of total change
Natural (environmental) factors		
The total surface water inflow into the lake	20	11
Precipitation at the lake surface	43	23
Evaporation from the lake water area	90	48
<i>Subtotal:</i>	<i>153</i>	<i>82</i>
Anthropogenic factors		
Runoff through the water interception drains	-24	-13
Water consumption e	-11	-6
The Sungach River runoff decrease due to reduction of its outflow from the Lake	35	19
Water inflow from the Small Khanka Lake (on the Chinese territory) due to the Muling River flood water diversions.	34	18
<i>Subtotal:</i>	<i>34</i>	<i>18</i>
TOTAL CHANGES	187	100