

## Submillimeter multi-wavelength observations for the protoplanetary disk around a young star TW Hya with ALMA

\*Takashi Tsukagoshi<sup>1</sup>, Hideko Nomura<sup>2</sup>, Takayuki Muto<sup>3</sup>, Ryohei Kawabe<sup>4,9,10</sup>, Daiki Ishimoto<sup>2,5</sup>, Kazuhiro Kanagawa<sup>6</sup>, Satoshi Okuzumi<sup>2</sup>, Shigeru Ida<sup>2</sup>, Catherine Walsh<sup>11,7</sup>, Tom Millar<sup>8</sup>

1. Ibaraki University, 2. Tokyo Institute of Technology, 3. Kogakuin University, 4. National Astronomical Observatory of Japan, 5. Kyoto University, 6. University of Szczecin, 7. Leiden University, 8. Queen's University Belfast, 9. The Graduate University for Advanced Studies, 10. University of Tokyo, 11. University of Leeds

We present the results of our  $\sim 3$  au resolution imaging observations of the protoplanetary disk around TW Hya at 145 and 233 GHz with the Atacama Large Millimeter/submillimeter Array. Our observations revealed two deep gaps at 22 and 37 au and shallower gaps at 6, 28, and 44 au, as reported by Andrews et al. (2016). The central hole with a radius of  $\sim 3$  au was also marginally resolved. The most remarkable finding is that the spectral index between bands 4 and 6 peaks at the 22 au gap. The derived power-law index of the dust opacity  $\beta$  is  $\sim 1.7$  at the 22 au gap and decreases toward the disk center to 0. The most prominent gap at 22 au could be caused by the gravitational interaction between the disk and an unseen Neptune-mass planet, although other origins may be possible. The planet-induced gap is supported by the fact that  $\beta$  is enhanced at the 22 au gap, indicating a deficit of millimeter-sized grains within the gap due to dust filtration by a planet.

Keywords: protoplanetary disk, planet formation, icy giant

