

Parameterized lattice strain model for HFSE partitioning between amphibole and silicate melt with application to arc magma evolution

*Kei Shimizu¹, Yan Liang², Erik H Hauri¹

1. Department of Terrestrial Magnetism, Carnegie Institution for Science, 2. Department of Earth, Environmental and Planetary Sciences, Brown University

The role of amphibole fractional crystallization in generating differentiation trends in arc magmas [1, 2] can be better understood by constraining the parameters responsible for the large range in experimentally determined trace element partition coefficients between amphibole and melt. In this study, we used published experimental high field strength element (HFSE) partitioning data between amphibole and silicate melt, the lattice strain model [3], and non-linear least squares regression method to parameterize key parameters in the lattice strain model (D_0 , r_0 , and E) for amphibole as a function of pressure, temperature, and both mineral and melt compositions. In our tetravalent HFSE (Zr, Hf, Ti) partitioning model, D_0 negatively correlates with Ti content in the melt, r_0 negatively correlates with Ti content in the amphibole, and E positively correlates with r_0 . The pentavalent HFSE (Nb, Ta) partition coefficients were empirically modeled as a function of amphibole composition and D_{HF} . Application of our HFSE partitioning models and our previous REE partitioning model [4] to experimental data simulating fractional crystallization of arc magmas [5] suggests that (1) HFSE and REE partition coefficients between amphibole and melt can vary by a factor of three and ten respectively during arc magma fractional crystallization due to variation in the amphibole and melt compositions, (2) amphibole fractional crystallization causes significant fractionation within and between HFSEs and REEs (e.g. it increases Zr/Hf and Zr/Nd ratios, and decreases Dy/Yb and Dy/Dy* ratios) in arc magmas, and finally (3) it plays a key role in buffering the HFSE and REE concentrations in arc magmas.

References

- [1] Davidson et al. (2007) *Geology* 35(9), 787-790.
- [2] Davidson et al. (2013) *J. Petrol.* 54(3), 525-537.
- [3] Blundy & Wood (1994) *Nature* 372, 452-454.
- [4] Shimizu et al., (2017) *Am. Mineral.* Submitted.
- [5] Nandedkar et al. (2014) *CMP* 167(6), 1-27.

Keywords: amphibole, HFSE and REE partitioning, arc magma evolution