

S-netデータによる東日本前弧域の構造と地震テクトニクス

Seismic structure and seismotectonics of the East Japan forearc derived from seafloor S-net data

*趙 大鵬¹、片山 悠¹、豊国 源知¹

*Dapeng ZHAO¹, Yu Katayama¹, Genti Toyokuni¹

1. 東北大学大学院理学研究科 地震・噴火予知研究観測センター

1. Research Center for Earthquakes & Volcanoes, Graduate School of Science, Tohoku University

A permanent seafloor seismic network (S-net) consisting of 150 stations has been recently installed in the East Japan forearc and the outer-rise area under the Pacific Ocean, which provides an unprecedented opportunity to investigate the detailed forearc structure and initial stage of plate subduction. Using P-wave arrival-time data of local suboceanic events recorded by the seafloor S-net, *Yu and Zhao* (2020) determined P-wave velocity (V_p) tomography off Hokkaido, *Hua et al.* (2020) obtained V_p tomography off Tohoku, and *Wang et al.* (2022) investigated 3-D V_p anisotropic structure off Tohoku. However, the 3-D S-wave velocity (V_s) structure of the forearc region has not been investigated using the S-net data, except for *Matsubara et al.* (2019). In this work, we use a large number of P and S wave arrival times of local earthquakes beneath the Pacific Ocean and land areas recorded by both the S-net and the land-based Kiban seismic network to investigate the 3-D V_p and V_s structures of the East Japan forearc. Geometries of the subducting Pacific slab boundary and the Conrad and Moho discontinuities under the forearc are refined by referring to results of previous studies including active-source seismic soundings and seismicity located by the S-net. The obtained V_p and V_s images are quite similar to each other, suggesting that they reflect robust features of the 3-D seismic structure. The subducting Pacific slab exhibits high velocities, whereas significant low-velocity (low- V) anomalies are revealed in the overlying Okhotsk plate beneath the forearc, which may reflect accretionary materials containing abundant fluids. The subducting oceanic crust atop the Pacific slab is revealed clearly as a thin low- V layer extending from the trench axis down to ~100 km depth under the volcanic front. Our results of earthquake relocation using the 3-D velocity model indicate that the double seismic zone in the subducting Pacific slab occurs from the outer-rise area east of the Japan trench down to ~180 km depth under the eastern margin of the Japan Sea. The occurrence and focal mechanisms of the double seismic zone are caused by slab deformation during its subduction, such as slab bending or unbending, as well as other processes including hydration, dehydration, compositional variations, and phase changes in the subducting slab. Significant lateral variations of velocity structure are also revealed in the interplate megathrust zone, which show correlations with the distribution of large interplate earthquakes. These results shed new light on the seismic structure and subduction dynamics of the forearc region (*Zhao et al.*, 2022).

References

- Hua, Y., D. Zhao, G. Toyokuni, Y. Xu (2020). Tomography of the source zone of the great 2011 Tohoku earthquake. *Nature Communications* 11, 1163.
- Matsubara, M., H. Sato, K. Uehira et al. (2019). Seismic velocity structure in and around the Japanese island arc derived from seismic tomography including NIED MOWLAS Hi-net and S-net data. *Seismic Waves - Probing Earth System*, IntechOpen, 1-18.
- Wang, Z.W., D. Zhao, X. Chen (2022). Seismic anisotropy and intraslab hydrated faults beneath the NE Japan forearc. *Geophys. Res. Lett.* 49, e2021GL097266.

- Yu, Z., D. Zhao (2020). Seismic evidence for water transportation in the forearc off Northern Japan. *J. Geophys. Res.* 125, e2019JB018600.
- Zhao, D., Y. Katayama, G. Toyokuni (2022). The Moho, slab and tomography of the East Japan forearc derived from seafloor S-net data. *Tectonophysics* 837, 229452.