

## オフセットダイポールを含む水星磁気圏における重イオンの運動論 Heavy ion kinetics in Mercury magnetosphere with offset dipole

八木 学<sup>1\*</sup>; 関 華奈子<sup>2</sup>; 松本 洋介<sup>3</sup>; Delcourt Dominique<sup>4</sup>; Leblanc Francois<sup>4</sup>  
YAGI, Manabu<sup>1\*</sup>; SEKI, Kanako<sup>2</sup>; MATSUMOTO, Yosuke<sup>3</sup>; DELCOURT, Dominique<sup>4</sup>; LEBLANC, Francois<sup>4</sup>

<sup>1</sup> 東北大学惑星プラズマ・大気研究センター, <sup>2</sup> 名古屋大学太陽地球環境研究所, <sup>3</sup> 千葉大学大学院理学研究科, <sup>4</sup> フランス国立科学研究センター

<sup>1</sup>PPARC, Tohoku University, <sup>2</sup>STEL, Nagoya University, <sup>3</sup>Chiba University, <sup>4</sup>CNRS

Based on observations by MESSENGER, Mercury magnetosphere is thought to be a miniature of the Earth magnetosphere. These two magnetospheres have several characteristics in common, however, some critical differences are also evident. First, there is no atmospheric layer, but only tenuous exosphere. Second, the kinetic effects of heavy ions might not be negligible because Mercury magnetosphere is relatively small compared to the large Larmor radii. Trajectory tracings is one of the dominant methods to estimate the kinetic effect of heavy ions which originate the exosphere, though the results of the simulation are quite sensitive to the electric and magnetic field. Therefore, it is important to provide a realistic field model in the trajectory tracings. In order to construct a large scale structure, we developed a MHD simulation code, and adopted to the global simulation of Mercury magnetosphere. We performed four solar wind conditions of the northward IMF, and the results showed that the global configurations such as the location of magnetopause depend heavily on the dynamic pressure, while the solar wind electric field contributes little to the magnetospheric configuration. On the other hand, the results of statistical trajectory tracings of exospheric sodium ions depend not only on the dynamic pressure but also on the solar wind electric field. In the results, we identified two efficient acceleration processes and formation of the sodium ring which is formed by the accelerated ions drifting around the planet by magnetic gradient of the dipole field. When the solar wind dynamic pressure is low, acceleration by magnetospheric convection is efficient in the vicinity of Mercury. When the dynamic pressure is high, entry of the accelerated ions picked-up in the magnetosheath into the magnetosphere becomes dominant. The entry point of sodium ions changes due to the variation of the solar wind electric field, which causes a difference in the sodium ring's shape for the same solar wind dynamic pressure cases. Recent observation by MESSENGER revealed the weaker dipole field of Mercury than the past estimation based on Mariner 10 as well as large offset of dipole which could change the global configuration of Mercury magnetosphere and behavior of sodium ions. In the presentation, we will also discuss the ongoing simulation including the above configuration of intrinsic magnetic field of Mercury especially focus on how will this affect the distribution of sodium ions and its acceleration mechanisms.

キーワード: 水星磁気圏, 磁気流体, テスト粒子, ナトリウムイオン, オフセットダイポール  
Keywords: Mercury magnetosphere, MHD, test particle, sodium ion, offset dipole