

SCG57-P17

会場:コンベンションホール

時間:5月27日 18:15-19:30

関東山地、上部白亜系跡倉層の碎屑性ジルコン年代学

Detrital zircon chronology of the Upper Cretaceous Atokura Formation in northern Kanto Mtn., Japan

中畠 浩基^{1*}; 磯崎 行雄²; 小坂 和夫³; 坂田 周平⁴; 平田 岳史⁴

NAKAHATA, Hiroki^{1*}; ISOZAKI, Yukio²; KOSAKA, Kazuo³; SAKATA, Shuhei⁴; HIRATA, Takafumi⁴

¹ 東京大学大学院理学系研究科地球惑星科学専攻, ² 東京大学大学院総合文化研究科広域科学専攻, ³ 日本大学文理学部地球システム科学科, ⁴ 京都大学大学院理学研究科地球惑星科学専攻

¹Department of Earth and Planetary Science, Graduate School of Science, The University of Tokyo, ²Department of Earth Science and Astronomy, The University of Tokyo, ³Department of Geosystem Sciences, College of Humanities and Sciences, Nihon University, ⁴Graduate school of Science, Kyoto University

The Cretaceous sandstones of the Atokura nappe in the northern Kanto Mtns. in central Japan were analyzed for detrital zircon U-Pb chronology by LA-ICP-MS. The results documented two distinct age groups; i.e., 150-300 Ma (Permian to Jurassic) and 1600-2600 Ma (Neoarchean to Mesoproterozoic). As there are no correlative candidates for provenance in the neighborhood, the Cretaceous strata of the Atokura nappe were likely exotic. The occurrence of Jurassic granitoids is currently limited to the Hida belt in central Japan, while no Precambrian basement is exposed at all in Japan. Judging from the allochthonous mode of occurrence, the Atokura nappe was derived originally from the back-arc domain of the Cretaceous Japan.

キーワード: 白亜紀, 砂岩, 碎屑性ジルコン, U-Pb 年代, LA-ICP-MS, 日本

Keywords: Cretaceous, sandstone, detrital zircon, U-Pb age, LA-ICP-MS, Japan