

Insight into the thermochemical state of the Jovian core from ab initio thermodynamic integration MD

TANIUCHI, Takashi^{1*} ; TSUCHIYA, Taku¹

¹Geodynamics Research Center, Ehime University

The thermochemical state of the Jovian rocky core still remains unclear. Although the rocky components of the cores in gas giants are likely to be MgO, SiO₂ and Fe (Guillot, 1999), reported melting condition are still limited up to ~500-1500 GPa (Boates and Bonev, 2013; Millot *et al.*, 2015; Bouchet *et al.*, 2013), which are far lower than the Jovian core P,T condition (4 TPa and 20 kK) estimated based on the equations of state of gas materials determined by ab initio calculations (Nettelmann *et al.*, 2012), primarily due to experimental difficulty. Thermochemical state (e.g., solid or liquid) of these materials might influence the tidal dissipation of planets and the mixing of the core and envelope. In order to clarify the phase equilibria, we have developed a new technique for calculating free energies of liquid and crystalline states based on the thermodynamic integration method (Frenkel and Smit, 2001) combined with the ab initio molecular dynamics method.

We obtained the following new findings: (1) calculated density of the MgO-SiO₂-Fe mixture is found highly consistent with the modeled Jovian core density, (2) although the melting temperatures of MgO and SiO₂ are higher than the modeled Jovian core temperature, the eutectic temperature is lower. These could serve for more detailed modeling of Jovian interior.

Keywords: ab initio calculation, thermodynamic integration method, Jovian core