

## Sources of atmospheric black carbon and related carbonaceous components at Rishiri Island, Japan

\*朱 春茂<sup>1</sup>、金谷 有剛<sup>1</sup>、吉川 久幸<sup>2</sup>、入野 智久<sup>2</sup>、関 幸<sup>2</sup>、遠嶋 康德<sup>3</sup>

\*Chunmao Zhu<sup>1</sup>, Yugo Kanaya<sup>1</sup>, Hisayuki Yoshikawa<sup>2</sup>, Tomohisa Irino<sup>2</sup>, Osamu Seki<sup>2</sup>, Yasunori Tohjima<sup>3</sup>

1. 海洋研究開発機構、2. 北海道大学、3. 国立環境研究所

1. Japan Agency for Marine-Earth Science and Technology, 2. Hokkaido University, 3. National Institute for Environmental Studies

A field study was conducted to clarify sources of atmospheric black carbon and related carbonaceous components at Rishiri Island, Japan. We quantified equivalent black carbon (eBC) particle mass and the absorption Ångström exponent (AAE), atmospheric CO and CH<sub>4</sub>, in addition to levoglucosan in total suspended particles, a typical tracer of biomass burning. Sixteen high eBC events were identified attributable to either anthropogenic sources or biomass burning in Siberia/China. These events were often accompanied by increases of co-emitted gases such as CH<sub>4</sub> and CO. Specifically, we observed pollution events with elevated eBC, AAE, levoglucosan, and CH<sub>4</sub>-CO slope in late July 2014, which were attributed to forest fires in Siberia by reference to the FLEXPART model footprint and fire hotspots. In autumn, drastic increases of eBC, AAE, and levoglucosan were observed, accompanied by an eBC-CO slope of >15 ng m<sup>-3</sup>/ppb, resulting from long-range transport of emissions from extensive burning of crop residue on the Northeast China Plain. Other than the sources of fossil fuel combustion in China and forest fires in Siberia, we also report that pollution events in northern Japan are caused by crop residue burning in China. This study elucidated valuable information that will improve understanding of the effects of biomass burning in East Asia on atmospheric carbonaceous components.

キーワード：エアロゾル、バイオマス燃焼、レボグルコサン、ブラックカーボン、FLEXPART

Keywords: Aerosols, Biomass burning, Levoglucosan, Black carbon, FLEXPART