

## 成層圏・対流圏オゾン交換の全球分布と経年変動の推定：気候場変動の影響

### Evaluating global distribution and interannual variation of stratosphere-troposphere exchange of ozone: impacts of climate variabilities

\*須藤 健悟<sup>1,2</sup>、Xin Li<sup>1</sup>

\*Kengo Sudo<sup>1,2</sup>, Xin Li<sup>1</sup>

1. 名古屋大学大学院環境学研究科、2. 海洋研究開発機構

1. Graduate School of Environmental Studies, Nagoya University, 2. Research and Development Center for Global Change, JAMSTEC

The stratosphere-troposphere exchange (STE) of ozone ( $O_3$ ) can have significant impacts on the interannual variability and long-term trend of global tropospheric ozone and radiation budget. This study investigates global STE of  $O_3$  particularly focusing on the roles of major climate variabilities such as ENSO, PDO, and QBO and climate trend (surface temperature). Using a chemistry coupled climate model, this study estimates temporal variation in the vertical and meridional STE  $O_3$  fluxes during the years of 2000-2016. For the major climate variabilities of ENSO and PDO, associated STE changes are assessed in detail using a composite analysis. The model validation shows that the model well captures the OMI-observed tropospheric column ozone (TCO) abundances ( $r \sim 0.55$ ), especially around the tropical Pacific ( $r \sim 0.8$ ). It is shown that in the tropics there is upward  $O_3$  transport to the stratosphere (troposphere to stratosphere transport, TST) while in the extratropics,  $O_3$  transport is downward (stratosphere to troposphere transport, STT), consistent with the Brewer-Dobson circulation in the stratosphere. Although geographical distributions of vertical and meridional STE fluxes are not changed largely by ENSO or PDO, the STE fluxes are significantly amplified by the ENSO events. Horizontal patterns of STE flux anomaly are found to be similar with respect to the El Niño episodes and the PDO warm phases (STT increased by 21% and 10%, respectively, in the northwest Pacific). These anomalies are reversed during the La Niña and PDO cold phases (STT decreased by 16% and 3%, respectively, in northeast Pacific). The anomaly of global net STE  $O_3$  flux is interpreted as several key factors using a multiple-regression analysis. The result shows that the STE anomaly is explained largely by the combination of ONI, PDO, QBO, and global mean surface temperature ( $R^2 \sim 43\%$ ). Contributions of ONI and PDO to STE anomaly are estimated to be 22.3% and 23.1%, respectively. This study quantitatively reveals the possible connection between climate condition and STE.

キーワード：成層圏・対流圏間交換、オゾン、化学気候モデル、化学気候相互作用、気候変動、対流圏オゾン収支

Keywords: Stratosphere-Troposphere Exchange (STE), Ozone, chemistry climate model, chemistry-climate interaction, climate change, Budget of tropospheric ozone