

Assessment of the effectiveness of the global ocean state estimation using tide-induced mixing parameterizations

*長船 哲史¹、増田 周平¹、杉浦 望実¹、土居 知将¹、邊見 忠¹

*Satoshi Osafune¹, Shuhei Masuda¹, Nozomi Sugiura¹, Doi Toshimasa¹, Tadashi Hemmi¹

1. 国立研究開発法人 海洋研究開発機構

1. JAMSTEC

Recent data synthesis experiments showed that adjusting mixing coefficients through data assimilation approach is a promising way to reduce a global misfit between a model simulation and ocean observations, while mixing coefficients should satisfy constraints from energetics. Aiming for a data synthesis experiment that is energetically consistent with the known constraint on the ocean energy budget and can assimilate turbulence observation data, we have developed a new quasi-global four-dimensional variational dataassimilation system using two parameterizations for tidally induced vertical mixing. As a preliminary preparation of the synthesis experiment, we optimally estimated their parameters along with other physical parameters based on the Green' s function method. The simulation using the optimal parameters well reproduces temperature and salinity in the deep Pacific Ocean. Using this simulation as the first guess field, we are conducting a long-term data synthesis experiment. We will present some preliminary results of the synthesis experiment to assess the effectiveness of our new system.

キーワード：鉛直混合、潮汐、データ同化

Keywords: vertical mixing, tide, data assimilation