

## Examination of $^{10}\text{Be}$ in travertine sediment as a possible tool to reconstruct past solar activity

\*Hongyang Xu<sup>1,2</sup>, Hiroko Miyahara<sup>2</sup>, Kazuho Horiuchi<sup>3</sup>, Hiroyuki Matsuzaki<sup>4</sup>, Hailong Sun<sup>5</sup>, Weijun Luo<sup>5</sup>, Xiangmin Zheng<sup>1</sup>, Yusuke Suganuma<sup>6,7</sup>, Shijie Wang<sup>5</sup>, Limin Zhou<sup>1</sup>

1. East China Normal University, 2. Musashino Art University, 3. Hirosaki University, 4. University of Tokyo, 5. Chinese Academy of Sciences, 6. National Institute of Polar Research, 7. The Graduate University for Advanced Studies (SOKENDAI)

In this study, we explore the possibility of using travertine sediment for reconstructing the past solar activity. We obtained an annual  $^{10}\text{Be}$  record from travertine deposit from the modern period and compared with the neutron monitor data. We discuss a possible methodology to calibrate the climatological impact on  $^{10}\text{Be}$  deposition and to obtain the signal of past solar and cosmic-ray variations.

Keywords: Beryllium-10, cosmogenic nuclide, solar activity, endogenic travertine