

## The model study of ionospheric variations during the solar eclipse event by assimilating the GNSS TEC observations

\*陳 佳宏<sup>1</sup>、Lin Charles<sup>1</sup>、Matsuo Tomoko<sup>2</sup>

\*Chia-Hung Chen<sup>1</sup>, Charles Lin<sup>1</sup>, Tomoko Matsuo<sup>2</sup>

1. 台湾国立成功大学地球科学学科、2. コロラド大学航空宇宙工学

1. Department of Earth Sciences, National Cheng Kung University, 2. Department of Aerospace Engineering Sciences, University of Colorado at Boulder, CO

Using the physics-based thermosphere-ionosphere model (NCAR-TIEGCM) with an ensemble Kalman filter, this study reports the first data assimilative analysis of the ionosphere responses to the solar eclipse on 21 August 2017. The system, using a 2-minute assimilation cycle of data from ground-based GNSS observations, show dynamic variations of the equatorial ionization anomaly (EIA) due to the electrodynamic effects of the solar eclipse. Two major ionospheric responses are captured: (1) an early appearance of EIA at the westward boundary of moon shadow and (2) an enhanced EIA at lower latitudes and suppressed EIA at the higher latitudes. These eclipse-induced conjugate EIA variations are produced by an eastward electric field perturbation around magnetic equator and a westward electric field perturbation at the higher latitudes.

キーワード：電離圏データ同化、日食、電場変動

Keywords: Ionospheric data assimilation, solar eclipse, electric field perturbation