

Correlations between plasma density and magnetic field strength in MHD turbulence in space

*羽田 亨¹、成田 康人²、成行 泰裕³

*Tohru Hada¹, Yasuhito Narita², Yasuhiro Nariyuki³

1. 九州大学大学院総合理工学研究院環境理工学部門、2. オーストリア科学アカデミー宇宙科学研究所、3. 富山大学人間発達科学部

1. Interdisciplinary Graduate School of Engineering Sciences, Kyushu University, 2. Space Research Institute, Austrian Academy of Sciences, 3. Faculty of Human Development, University of Toyama

Large amplitude magnetohydrodynamic (MHD) turbulence is ubiquitous in space plasma, for example, in the solar wind, in the foreshock region (Narita, World Scientific, 2010) and in the sheath region of the earth's bowshock (e.g., Pollock et al., J. Atmosph. Solar Terr. Phys, 2018). Moreover, recent Voyager observations of large amplitude density and magnetic field fluctuations in the heliosheath behind the termination shock arouse a vivid discussion on the compressible nature and the origin of the observed turbulence (Burlaga and Ness, ApJ, 2009; Gutynska et al., ApJ, 2010). An obvious way of generating the density fluctuations is via the presence of compressive wave modes, such as obliquely propagating MHD waves. The density response is linear to the magnetic field perturbations in this case, and it vanishes in the limit of parallel wave propagation. In a finite (large) amplitude Alfvénic turbulence the density perturbation may be mainly generated via the "frozen-in" of the magnetic field to the plasma or the static balance between the plasma pressure and the magnetic field. Both of these processes result in the quadratic response of the density to the magnetic field perturbations, although their correlation is positive for the former and negative for the latter. This simple fluid picture, sometimes referred to as the "quasi-static" approximation, is known to be significantly modified when the ion response to the turbulence is kinetically treated (Mjølhus and Wyller, Phys. Scripta, 1986; Medvedev and Diamond, Phys. Plasmas, 1996). In this presentation, we address this unsolved issue of the correlations between the plasma density and the magnetic field amplitude in the presence of finite amplitude MHD turbulence by performing hybrid simulations (kinetic ions + fluid electrons), and by examining the in situ Cluster spacecraft data of the foreshock plasma (Narita and Hada, Earth, Planets and Space, 2018).

キーワード：MHD 乱流、準静的条件

Keywords: MHD turbulence, quasi-static approximation