

Ocean bottom seismographic observation and similar earthquake detection in the northern part of the Nansei Islands

*仲谷 幸浩¹、八木原 寛¹、平野 舟一郎¹、小林 励司²、宮町 宏樹²、中尾 茂²、内田 和也³、松島 健³、清水 洋³、中東 和夫⁴、山下 裕亮⁵、阿部 英二⁶、山田 知朗⁶、篠原 雅尚⁶

*Yukihiro Nakatani¹, Hiroshi Yakiwara¹, Shuichiro Hirano¹, Reiji Kobayashi², Hiroki Miyamachi², Shigeru Nakao², Kazunari Uchida³, Takeshi Matsushima³, Hiroshi Shimizu³, Kazuo Nakahigashi⁴, Yusuke Yamashita⁵, Hideji Abe⁶, Tomoaki Yamada⁶, Masanao Shinohara⁶

1. 鹿児島大学附属南西島弧地震火山観測所、2. 鹿児島大学大学院理工学研究科、3. 九州大学大学院理学研究院附属地震火山観測研究センター、4. 東京海洋大学、5. 京都大学防災研究所、6. 東京大学地震研究所

1. Nansei-Toko Observatory for Earthquakes and Volcanoes, Kagoshima University, 2. Graduate School of Science and Engineering, Kagoshima University, 3. Institute of Seismology and Volcanology, Faculty of Science, Kyushu University, 4. Tokyo University of Marine Science and Technology, 5. Disaster Prevention Research Institute, Kyoto University, 6. Earthquake Research Institute, the University of Tokyo

The Ryukyu Trench subduction zone is seismically active region. In the northern part of the Nansei Islands, both seismic and aseismic events have occurred along the subducting plate interface. Spatial distribution of small local earthquakes in this region seems to show some seismic clusters and gaps. Short-term slow slip events and shallow low-frequency tremors have been also observed by recent onshore seismic and geodetic data. However, patterns of their slip behavior and spatiotemporal relationship between regular and slow earthquakes are less well understood because seismic stations are located only at the islands that are linearly arranged; at least 100 km away from the Ryukyu Trench. We have therefore conducted repeating long-term ocean bottom seismographic observation off the northern part of the Nansei Islands since 2014. First, we deployed three pop-up type ocean bottom seismometers (OBSs) in April 2014. Then, three more OBSs were added to the network in July 2015, and after that deployment and retrieval of OBSs at the same position have been repeatedly carried out every about one year. The station interval is about 40–60 km.

In this study, we focus on regular local earthquakes, particularly seismic clusters beneath the central part of the observation network. In order to obtain key properties of the clusters and to detect missing earthquakes, we apply a matched-filter technique to OBS continuous waveform data. As template event waveforms, we select events determined by Nansei-Toko Observatory for Earthquakes and Volcanoes, Kagoshima University. The hypocenters of the template events are relocated by merging manual pick data of P phases and S phases recorded on the OBSs with those on the onshore stations. As a preliminary result, we achieve detection of missing earthquakes using example template events and a threshold rule based on the median absolute deviation of the mean correlation coefficient. In this presentation, we further discuss the characteristics of waveform similarity of the OBS data.

Acknowledgements: This work is supported by Earthquake and Volcano Hazards Observation and Research Program, the MEXT of Japan and by ERI JURP.

キーワード : OBS observation, the Ryukyu Trench, similar earthquakes

Keywords: OBS observation, the Ryukyu Trench, similar earthquakes