

Methane emission estimate from South Asia: AMASA project

Methane emission estimate from South Asia: AMASA project

*林田 佐智子¹、Chandra Naveen¹、Patra Prabir²、寺尾 有希夫³、須藤 重人⁴、犬伏 和之⁵、間野 正美⁵、山本 昭範⁶

*Sachiko Hayashida¹, Naveen Chandra¹, Prabir Patra², Yukio Terao³, Shigeto Sudo⁴, Kazuyuki Inubushi⁵, Masayoshi Mano⁵, Akinori Yamamoto⁶

1. 奈良女子大学理学部、2. 海洋研究開発機構、3. 国立環境研究所、4. 農業環境技術研究所、5. 千葉大学大学院園芸研究科、6. 東京学芸大学

1. Faculty of Science, Nara Women's University, 2. Japan Agency for Marine-Earth Science and Technology (JAMSTEC), 3. National Institute for Environmental Studies (NIES), 4. National Institute for Agro-Environmental Sciences (NIAES), 5. Graduate School of Horticulture, Chiba University, 6. Tokyo Gakugei University

Methane (CH₄) is the second most significant anthropogenic greenhouse gas (IPCC AR5, 2014) after carbon dioxide (CO₂). Large fraction of CH₄ emissions (~150 Tg of 550 Tg globally; 1Tg=10¹²g) occur in the Asia region. Much of CH₄ emissions from Asia are attributable to ruminant animals and rice fields, but the quantitative estimate of those emissions remains highly uncertain. To improve CH₄ emission estimate from South Asia, we started a project “Atmospheric Methane and Agriculture in South Asia (AMASA)”, which is sponsored by the Ministry of the Environment Japan. The first goal of the project is to develop high-resolution emission maps at regional scale and improve our understanding of CH₄ emission distributions from South Asia by using remote sensing data of CH₄ from Japanese satellite Greenhouse Gases Observing Satellite (GOSAT), in-situ measurements at ground-based stations and atmospheric chemistry-transport model (ACTM) simulations. The second goal is to develop an emission mitigation proposal using results from India-specific rice field experiments for different management practices. Some emission mitigation scenarios will be developed based on the field data, and using the ACTM we examine to what extent the emission reductions are detectable by the measurement systems if the emission mitigation policy is realized.

GOSAT is the first satellite that is dedicated to greenhouse-gas-monitoring. The onboard Sensor TANSO-FTS (Fourier Transform Spectrometer) is designed to measure CO₂ and CH₄. GOSAT has collected data for about 8 years, and validation studies revealed sufficient reliability of GOSAT data for CH₄ cycling in the Earth's environment (e.g., Morino et al., 2011, Inoue et al., 2014, Ono et al., 2015). The essential merit of satellite observation is wide spatial coverage. We found very high concentrations of CH₄ over Asia in GOSAT data, which seem to be connected to the high CH₄ emissions from this region. However, the connection is not straightforward because of complicated transport mechanisms. In particular, at the foothills of Himalaya Mountains, upwelling wind lift the CH₄ up to mid- and upper-troposphere during the monsoon season, resulting in the high columnar concentrations that can be observed from GOSAT (Chandra et al., 2017, paper in preparation). We are also conducting field measurements of atmospheric CH₄ at ground-based stations in Karnal, Sonapat, Nainital, and Comilla. Combined use of those satellite-based and ground-based measurements can give us spatial structure of CH₄ distribution, which would improve the emission estimate using an inverse analysis system.

To develop an emission mitigation proposal, we are conducting rice field experiments at the Tamilnadu Rice Research Institute, India by managing rice cultivars, water table and soil properties. The preliminary results suggest that CH₄ emission from rice cultivation can be reduced by half when applying proper cultivation managements. Based on these new findings, we are pursuing an appropriate mitigation proposal to reduce CH₄ emissions from South Asia.

References

Ono, A. et al., Comparison of GOSAT SWIR and Aircraft Measurements of XCH_4 over West Siberia, SOLA, 11, 160-164, doi:10.2151/sola.2015-036, 2015.

Inoue, M. et al., Validation of XCH_4 derived from SWIR spectra of GOSAT TANSO-FTS with aircraft measurement data, Atmos. Meas. Tech., 7, 2987-3005, doi:10.5194/amt-7-2987-2014, 2014.

Morino, I. et al.: Preliminary validation of column-averaged volume mixing ratios of carbon dioxide and methane retrieved from GOSAT short-wavelength infrared spectra, Atmos. Meas. Tech., 4, 1061-1076, doi:10.5194/amt-4-1061-2011, 2011.

キーワード：メタン、温室効果気体、アジア

Keywords: methane, greenhouse gas, Asia

