

Higher temperature accelerate the growth of iron-limited phytoplankton communities in the subarctic Pacific

*杉江 恒二¹、芳村 毅²、西岡 純³

*Koji Sugie¹, Takeshi Yoshimura², Jun Nishioka³

1. 国立研究開発法人海洋研究開発機構 地球環境観測研究開発センター、2. 一般財団法人電力中央研究所 環境科学研究所、3. 国立大学法人北海道大学低温科学研究所 環オホーツク観測研究センター

1. Research & Development Center for Global Change, Japan Agency for Marine-Earth Science and Technology, 2. Environmental Science Research Laboratory, Central Research Institute of Electric Power Industry, 3. Pan-Okhotsk Research Center, Institute of Low Temperature Science, Hokkaido University, Hokkaido University

Iron is one of the key element determining the carrying capacity of primary production in the marine ecosystem. Because of iron scarcity in certain oceanic regions, phytoplankton communities were dominated by small-sized non-diatom species and thus less efficient trophic transfer and biological carbon pump. However, the effect of climate change such as warming on iron-limited phytoplankton is poorly investigated. This study investigated that the effect of higher temperature (+ 4–5°C relative to controls) and iron addition on the growth dynamics of phytoplankton communities in the subarctic Pacific. Two experiments were conducted during summer in 2014 aboard R/V *Mirai* (MR14W: 47°N, 160°E; MR14E: 47°N, 148°W) and another one experiment was conducted during late winter in 2015 aboard R/V *Hakuho-Maru* (KH15: 42°N, 146°E). The addition of iron enhanced the specific growth rate of larger phytoplankton groups in all three experiments whereas smaller phytoplankton groups were rarely affected. In MR14W and KH15 experiments, higher temperature enhanced the growth of phytoplankton in both unamended controls and iron-added treatments. The magnitude of growth enhancement by the temperature increases was larger in smaller phytoplankton groups relative to the larger groups in the KH15 experiment. According to the relative growth rate analysis (between higher temperature (+ 4–5°C) treatment to control temperature under each iron conditions, and between iron-added to non-iron-added treatment under each temperature conditions), we found that the summer phytoplankton communities were iron-limited rather than temperature-limited whereas late winter phytoplankton community was temperature-limited rather than iron-limited. Our results suggest that the temperature conditions could modify the dynamics of iron-limited phytoplankton communities, especially for the low-temperature conditions. In addition, future global warming may enhance the growth of phytoplankton even in the iron-limited open ocean. Given the higher temperature could enhance the growth of smaller phytoplankton groups relative to the larger ones as observed in the KH15 experiment, not only climate change but also seasonal warming from winter to summer may play a key role in the dynamics of community composition in the ocean.

キーワード：温暖化、植物プランクトン、亜寒帯北太平洋

Keywords: Global warming, phytoplankton, subarctic Pacific Ocean