

## Response of sea surface fugacity of CO<sub>2</sub> to the Southern Annular Mode (SAM) shift south of Tasmania

\*Liang Xue<sup>1</sup>, Libao Gao<sup>1</sup>, Weidong Yu<sup>1</sup>, Meng Wei<sup>1</sup>

1. FIO, SOA, China

Using observational data collected south of Tasmania during 14 austral summer cruises during 1993-2011, we examined the response of sea surface fugacity of carbon dioxide ( $f\text{CO}_2$ ) to the Southern Annular Mode (SAM) shift, which occurred around 2000. In the southern part of the Southern Ocean (SO) or the Polar Zone (PZ) and the Polar Frontal Zone (PFZ),  $f\text{CO}_2$  increased faster at the sea surface than in the atmosphere before the SAM shift, but not after the shift. In the northern part of the SO or the Sub-Antarctic Zone (SAZ), however, surface  $f\text{CO}_2$  increased faster than atmospheric  $f\text{CO}_2$  both before and after the shift. The SAM shift had an important influence on the surface  $f\text{CO}_2$  trend in the PZ and PFZ, but not in the SAZ, which we attribute to differences in regional oceanographic processes (upwelling vs. non-upwelling). The SAM shift may have reversed the negative trend of SO CO<sub>2</sub> uptake.

Keywords: Southern Ocean, Southern Annular Mode shift, Carbon cycling