

Lithology and Depositional Age of Paleoproterozoic Volcaniclastic Sequence from Ashanti Belt of Birimian Supergroup, Southwest Ghana

*吉丸 慧¹、清川 昌一¹、坂井 志緒乃¹、伊藤 孝²、池原 実³、堀江 憲路⁴、竹原 真美⁴、佐野 貴司⁵、Nyame Frank⁶、Tetteh Georg⁷

*Satoshi Yoshimaru¹, Shoichi Kiyokawa¹, Shiono Sakai¹, Takashi Ito², Minoru Ikehara³, Kenji Horie⁴, Mami Takehara⁴, Takashi Sano⁵, Frank K Nyame⁶, George M Tetteh⁷

1. 九州大学理学府地球惑星科学専攻、2. 茨城大学、3. 高知大学、4. 国立極地研究所、5. 国立科学博物館、6. ガーナ大学、7. タルクワ鉱山技術大学

1. Department of Earth and Planetary Science, Kyushu University, 2. Ibaraki University, 3. Kochi University, 4. National Institute of Polar Research, 5. National Museum of Nature and Science, 6. University of Ghana, 7. University of Mines and Technology

Paleoproterozoic Birimian volcaniclastic successions occur along the coast near Cape Three Points, in Ashanti belt, southwest Ghana. In this study, the depositional environments and bioactivities recorded in the c.a. 2.3 Ga rocks were investigated. We report the structure and stratigraphy in the area, improved depositional age of the sediments by zircon U-Pb dating and C isotope features of the bioactivities in the paleo-ocean.

The Ashanti belt, generally showing NE-SW strike, composed mainly of andesitic basalts, volcaniclastic rocks and belt type granitoids (Perrouy et al., 2012). The greenstone is unconformably overlain by Tarkwaian conglomerates and metasedimentary rocks. The maximum depositional age of the overlying metasedimentary rocks and the oldest age of the intruded rock into Birimian volcanics in the Sekondi region is 2154 ± 2 Ma and 2174 ± 2 Ma, respectively (U-Pb zircon: Oberthür et al., 1998).

Detailed field investigations were performed to reconstruct about 1000m out of over 2000m thick stratigraphy of the volcaniclastic sequence in the Cape Three Points area. The rocks, affected by greenschist-facies metamorphism, generally strike N-S, mainly dip 60-80° to the east and show fining upward stratigraphy. Using a west vergent thrust fault which occurs in the central part of the area, we sub-divided the area into two, a western zone and an eastern zone. The eastern zone consists of approximately 800 m thick fining upwards sequence which appears to have been repeated two times within the sequence. Thick basaltic lava in the lowermost part of the sequence is replaced by or changes into sediments produced by gravity flows. The uppermost part of the sequence reveals thick basaltic andesite and/or pyroclastic layers which then changes to fine altered volcaniclastics containing organic material. The western zone has not yet been reconstructed the stratigraphy.

TiO₂/Al₂O₃ ratios from EPMA analyses of chromite in basaltic rocks suggest that these rocks originated in a volcanic arc system. Whole-rock trace element compositions tend to show low concentrations of Nb and high LREEs, which also supports derivation from volcanic arc.

A foliated porphyritic dyke which occurs in the Cape Three Points area was dated by SHRIMP at the National Institute of Polar Research. Zircon grains in the dyke yielded a weighted mean ²⁰⁴Pb-corrected ²⁰⁷Pb/²⁰⁶Pb age of 2265.6 ± 4.6 Ma (95% confidence), which suggests that the volcaniclastic sequence was deposited before 2265.6 ± 4.6 Ma and was deformed afterwards. This age is the oldest in the Ghanaian greenstone terrane (Loh and Hirdes, 1999). It would seem, therefore, that rocks in the Cape Three Points area record the history of early volcanic activity in the Birimian greenstone terrain.

We measured organic carbon isotope ratio of some black shale samples. The TOC varies from 0.02 %- 0.3 % and the δ¹³C values are -35 ‰ - -15 ‰ (N=5). These low concentrations of the organic matter suggest dilution by the continual influx of volcanogenic clastics.

The thick fining-upward volcaniclastic sequence and chemical compositions may suggest the Cape

Three Points Formation was deposited on mid-deep sea floor beside an oceanic volcanic arc. The deposition may have occurred before 2265.6 ± 4.6 Ma suggesting these sediments record oceanic environmental changes from Makganyene glaciation toward Lomagundi Event (2.3 Ga -2.06 Ga, Bekker and Holland, 2012 and Kopp et al., 2005).

キーワード：ガーナ・ビリミアン超層群、古原生代

Keywords: Birimian Supergroup, Ghana, Paleoproterozoic