

東エジプト El Dabbah 地域の層序と堆積場の復元：新原生代の縞状鉄鉱層 Reconstruction of stratigraphy and tectonic setting in El Dabbah, Eastern Egypt: Neoproterozoic banded iron formations

*鈴木 大志¹、清川 昌一¹、池原 実²、佐野 貴司³、ダウッド マヘル⁴、アブエルハッサン モハメッド⁴
*TAISHI SUZUKI¹, Shoichi Kiyokawa¹, Minoru Ikehara², Takashi Sano³, Maher Dawood⁴,
Mohamed Abouelhassan⁴

1. 九州大学、2. 高知大学、3. 国立科学博物館、4. メノフィア大学

1. kyushu Univ., 2. Kochi Univ., 3. National Museum of Nature and Science, 4. Menoufia Univ.

In the Eastern Desert of Egypt, Neoproterozoic Banded Iron Formations (BIFs) are reported within Nubia greenstone belt whose geological structure and stratigraphy are not well understood. In this study, the geological structure was established to reconstruct the stratigraphy including Iron formations (IFs) in El Dabbah in the middle of the Eastern Desert.

Greenstone sequence in El Dabbah area, strike-slip basin on Nubia shield during pan African orogeny, is covered by Hammamat Group. The greenstone sequence preserved the thick volcano-sedimentary rocks with gabbro, coarse-grained volcanoclastics rocks, pillow lavas, black shales, and BIFs. Reconstruction of stratigraphy in this greenstone sequence, which is 4000m in total thickness, composes Basal, Lower, Middle, and Upper members. There are 13 iron sections within this greenstone sequence.

Especially, BIFs sequence preserved within Lower and Upper members. BIFs sequences are well preserved within massive - pillow lavas sections and it contain laminated greenish - black shales. BIFs are composed of magnetite and/or hematite. Magnetite are oxidized from rims and become hematite.

We have determined major and trace element compositions of whole rock of volcanics and BIFs by using XRF and ICP-MS analysis. Trace elements compositions indicate that the volcanics are arc origin in term of a Nb-Zr-Y discrimination diagram and a MORB normalized spider diagram. Major elements in BIFs in this area are characterized by their higher content of Al_2O_3 than the other Archean BIFs. In addition, REE in BIFs show pattern of light REE enrichment and negative Eu anomalies.

This Volcanoclastic sedimentary sequence would be related at island arc sedimentary sequence with volcanic activity.

キーワード：縞状鉄鉱層、新原生代

Keywords: banded iron formation, Neoproterozoic