

Paleoceanographic proxy and skeletal records of sclerosponges from the Ryukyu Islands, Japan

*浅海 竜司¹、松森 建人¹、石原 信司¹、金城 章¹、大城 大輝¹、成瀬 貫¹、水山 克¹、伊勢 優史²、藤田 喜久³、坂巻 隆史⁴

*Ryuji Asami¹, Taketo Matsumori¹, Shinji Ishihara¹, Akira Kinjo¹, Daiki Ohshiro¹, Tohru Naruse¹, Masaru Mizuyama¹, Yuji Ise², Yoshihisa Fujita³, Takashi Sakamaki⁴

1. 琉球大学、2. 名古屋大学、3. 沖縄県立芸術大学、4. 東北大学

1. University of the Ryukyus, 2. Nagoya University, 3. Okinawa Prefectural University of Arts, 4. Tohoku University

Hypercalcified sponges (hereafter, “sclerosponges”), living in dark environments of tropical to subtropical shallow oceans, precipitate calcium carbonate skeleton with growth bands. They grow slowly at an approximate rate of <1 mm/year unlike corals (ca. 1 cm/year) but can be so long-lived for several decades to hundred years like corals (e.g., Benavides and Druffel, 1986). Skeletal oxygen isotopic ratios in sclerosponges can reflect variations in sea surface temperature and seawater with the latter being potentially related to salinity reflecting the precipitation–evaporation balance at the sea surface and changes in water mass transport (e.g., Grottoli et al., 2010). In contrast to zooxanthellate corals, which occasionally show positive correlations between skeletal oxygen and carbon isotopic ratios, it is considered that there do not exist vital effects in the secretion of sclerosponge skeleton. Previous studies showed significant decrease trends in the carbon isotope records toward the present, which is probably a result of isotopically-light carbon dioxide added into the atmosphere/ocean from fossil fuel burning (e.g., Swart et al., 2010). Therefore, sclerosponges are shown to provide annually resolved time series of proxy records of ocean environments since the Industrial Revolution. However, few evaluation studies on the environmental proxy of sclerosponges are reported and temporally longer (>100 year-continuous-long) records from sclerosponges were derived only from the Atlantic Ocean.

Here we present oxygen and carbon isotope records from sclerosponges (*Astrosclera willeyana* and *Acanthochaetetes wellsii*) collected from Kume-jima, Okinawa-jima and Miyako-jima, Ryukyu Islands in the North Pacific. In order to evaluate the utility of sclerosponge as an environmental proxy, within-skeleton and intraspecific variations in stable isotopic records of 37 samples were investigated. Soft X-ray images of large-size sclerosponges showed highly developed skeletal growth bands with >100 high/low density layers. The secular changes in radiocarbon-dated time series of the sclerosponge carbon records were consistent with previously reported data from Atlantic and Pacific corals and sclerosponges. The long-term oxygen isotopic trends of the samples are characterized by slight depletions throughout their living periods, indicative of an overall trend toward warmer ocean environment around the Ryukyu Islands. Our sclerosponge-based estimates of sea surface temperature and salinity may document thermal and hydrologic variations in the Ryukyu Islands, furthering a better understanding of northwestern tropical-subtropical Pacific climate change for the last several centuries in conjunction with coral-based long proxy records.

Benavides and Druffel (1986) Coral Reefs, 4, 221-224.

Grottoli et al. (2010) J. Geophys. Res., 115, doi:10.1029/2009JC005586.

Swart et al. (2010) Geophys. Res. Lett., 37, doi:10.1029/2009GL041397.

キーワード：硬骨海綿、骨格、プロキシ、酸素同位体組成、炭素同位体組成、古海洋

Keywords: sclerosponge, skeleton, proxy, oxygen isotope composition, carbon isotope composition, paleocean

